

Product datasheet

info@arigobio.com

ARG22075 anti-NK1.1 / CD161bc antibody [PK136] (low endotoxin)

Package: 250 μg Store at: -20°C

Summary

Product Description Azide free and low endotoxin Mouse Monoclonal antibody [PK136] recognizes NK1.1 / CD161bc

Tested Reactivity Ms

Tested Application BL, Cell-Act , Depletion, FACS, IHC, IP

Specificity Mouse NK1.1

Host Mouse

Clonality Monoclonal

Clone PK136

Isotype IgG2a, kappa

Target Name NK1.1 / CD161bc

Species Mouse

Immunogen CE Mouse spleen enriched for NK-1+ cells and bone marrow cells

Conjugation Un-conjugated

Alternate Names CD antigen CD161c; Al462337; Ly-55c; Nkrp1c; NK-RP1; NKR-P1C; Ly-59; Killer cell lectin-like receptor

subfamily B member 1C; NKR-P1.9; Ly55c; NKRP1; NKR-P1 40; ly-55c; Ly59; Nk1.2; NKRP140; CD161 antigen-like family member C; CD161; Lymphocyte antigen 55c; Nk1; Natural killer cell surface protein

P1-40; Nk-1.2; NK1.1; Nk-1

Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Cell-Act	Assay-dependent
	Depletion	Assay-dependent
	FACS	< 1 µg/10^6 cells
	IHC	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification Note	Low endotoxin
Buffer	PBS

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links <u>GeneID: 17059 Mouse</u>

Swiss-port # P27814 Mouse

Gene Symbol Kirb1c

Gene Full Name killer cell lectin-like receptor subfamily B member 1C

Function Plays a stimulatory role on natural killer (NK) cells cytotoxicity. Activation by cross-linking of the

receptor induces Ca(2+) mobilization and interferon-gamma production. [UniProt]

Calculated Mw 25 kDa