

**ARG22588**  
**anti-CLEC2 / CLEC1B antibody [17D9]**Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Rat Monoclonal antibody [17D9] recognizes CLEC2 / CLEC1B This antibody recognizes mouse C-type lectin domain family 1 member B, also known as CLEC2 or CLEC1B. CLEC2 acts as a receptor for rhodocytin, a platelet-aggregating protein found in snake venom. CLEC2 expression is not restricted to platelets and also acts as an activation receptor on neutrophils.
Tested Reactivity	Ms
Tested Application	FACS, WB
Host	Rat
Clonality	Monoclonal
Clone	17D9
Isotype	IgG2b
Target Name	CLEC2 / CLEC1B
Species	Mouse
Immunogen	RBL-2H3 cells stably expressing HA-tagged mouse CLEC2.
Conjugation	Un-conjugated
Alternate Names	CLEC2; C-type lectin domain family 1 member B; CLEC2B; PRO1384; C-type lectin-like receptor 2; CLEC-2; 1810061I13Rik; QDED721

### Application Instructions

Application table	Application	Dilution
	FACS	Neat
	WB	Assay-dependent
Application Note	FACS: Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% Sodium azide
Preservative	0.09% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

**Note**

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Gene Symbol	Clec1b
Gene Full Name	C-type lectin domain family 1, member b
Background	Natural killer (NK) cells express multiple calcium-dependent (C-type) lectin-like receptors, such as CD94 (KLRD1; MIM 602894) and NKG2D (KLRC4; MIM 602893), that interact with major histocompatibility complex class I molecules and either inhibit or activate cytotoxicity and cytokine secretion. CLEC2 is a C-type lectin-like receptor expressed in myeloid cells and NK cells (Colonna et al., 2000 [PubMed 10671229]).[supplied by OMIM, Jan 2011]
Calculated Mw	27 kDa
PTM	Glycosylated. Phosphorylated on tyrosine residue in response to rhodocytin binding.