

Product datasheet

info@arigobio.com

ARG22637 anti-O-GlcNAc antibody [9H6]

Package: 50 μg Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody [9H6] recognizes O-GlcNAc

Tested Reactivity Other

Tested Application ICC/IF, WB

Specificity Specific for β-O-Linked N-Acetylglucosamine modified proteins. Does not cross-react with N-

Acetylgalactosamine, Glucose, or Galactose modified proteins.

Host Mouse

Clonality Monoclonal

Clone 9H6

Isotype IgM

Target Name O-GlcNAc
Species Others

Immunogen Synthetic β-O-Linked N-Acetylglucosamine conjugated to KLH.

Conjugation Un-conjugated

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|----------|
| | ICC/IF | 1:50 |
| | WB | 1:1000 |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

Form Liquid

Purification PEG Purified.

Buffer PBS (pH 7.4), 0.9% Sodium azide and 50% Glycerol.


Preservative 0.9% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot


and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.



ARG22637 anti-O-GlcNAc antibody [9H6] ICC/IF image

Immunofluorescence: 5% Formaldehyde fixed (5 min) HEK293 cells stained with ARG22637 anti-O-GlcNAc antibody [9H6] at 1:50 for 30 - 60 min at RT. Secondary Antibody: Goat anti-Mouse (Alexa Fluor 488) at 1:1500 for 30 - 60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain. (A) DAPI (blue) nuclear stain. (B) Phalloidin Alex Fluor 633 F-Actin stain. (C) Primary antibody (D) Composite. .



ARG22637 anti-O-GlcNAc antibody [9H6] WB image

Western blot: 2.0 μ g of BSA, GlcNAc-BSA, GalNAc-BSA, Galactose-BSA, Glucose-BSA, and Citrulline-BSA stained with ARG22637 anti-O-GlcNAc antibody [9H6] at 1:1000 for 2 hours at RT.