

ARG23234
anti-MUC2 / Mucin 2 antibody [996/1]Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [996/1] recognizes MUC2 / Mucin 2
Tested Reactivity	Hu
Tested Application	FACS, IHC-Fr, IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	996/1
Isotype	IgG1
Target Name	MUC2 / Mucin 2
Species	Human
Immunogen	MUC2 tandem repeat peptide.
Conjugation	Un-conjugated
Alternate Names	MUC-2; MLP; Intestinal mucin-2; SMUC; Mucin-2

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	Assay-dependent
	WB	Assay-dependent

Application Note FACS: Membrane permeabilization is required for this application. Use 10 µl of the suggested working dilution to label 10⁶ cells in 100 µl.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% Sodium azide.
Preservative	0.09% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	MUC2
Gene Full Name	mucin 2, oligomeric mucus/gel-forming
Background	<p>This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. The protein polymerizes into a gel of which 80% is composed of oligosaccharide side chains by weight. The protein features a central domain containing tandem repeats rich in threonine and proline that varies between 50 and 115 copies in different individuals. Alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known. [provided by RefSeq, Jul 2008]</p>
Function	<p>Coats the epithelia of the intestines, airways, and other mucus membrane-containing organs. Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces. Major constituent of both the inner and outer mucus layers of the colon and may play a role in excluding bacteria from the inner mucus layer. [UniProt]</p>
Calculated Mw	540 kDa
PTM	<p>O-glycosylated.</p> <p>May undergo proteolytic cleavage in the outer mucus layer of the colon, contributing to the expanded volume and loose nature of this layer which allows for bacterial colonization in contrast to the inner mucus layer which is dense and devoid of bacteria.</p> <p>At low pH of 6 and under, undergoes autocatalytic cleavage in vitro in the N-terminal region of the fourth VWD domain. It is likely that this also occurs in vivo and is triggered by the low pH of the late secretory pathway. [UniProt]</p>