

Product datasheet

info@arigobio.com

ARG23778 Goat anti-Mouse IgA + IgG + IgM antibody (AP), pre-adsorbed

Package: 500 μl Store at: 4°C

Summary

Product Description AP-conjugated Goat Polyclonal antibody recognizes Mouse IgA + IgG + IgM

Tested Reactivity Ms

Tested Application ELISA, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB

Specificity Reacts with the heavy and light chains of mouse IgM, IgG1, IgG2a, IgG2b, IgG3 and IgA as demonstrated

by ELISA, FLISA, and/or flow cytometry; minimal cross reactivity with human immunoglobulins

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name IgA + IgG + IgM

Species Mouse
Conjugation AP

Application Instructions

Pre Adsorbed	Human immunoglobulins and pooled se	ra.
--------------	-------------------------------------	-----

Application table Application Dilution

Application	Dilution
ELISA	1:2000 - 1:4000
FACS	Assay-dependent
FLISA	Assay-dependent
ICC/IF	Assay-dependent
IHC-Fr	Assay-dependent
IHC-P	Assay-dependent
WB	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Properties

Form Liquid

Purification Affinity purification with immunogen.

Buffer Tris (pH 8.0), 0.1% Sodium azide, 1 mM MgCl2 and 50% Glycerol.

Preservative 0.1% Sodium azide

Stabilizer 50% Glycerol

www.arigobio.com argo.nuts about antibodies 1/2


Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Images



ARG23778 Goat anti-Mouse IgA + IgG + IgM antibody (AP) (preadsorbed) IHC-P image

Immunohistochemistry: Paraffin-embedded chicken nasal cavity section infected with H7N1 highly pathogenic avian influenza virus was stained with anti-IAV NP followed by ARG23778 Goat anti-Mouse IgA + IgG + IgM antibody (AP) (pre-adsorbed) and BCIP/NBT and then anti-PGP9.5 followed a secondary antibody and DAB.