

ARG23989 anti-Collagen III antibody [8D1-8C7]

Package: 50 µl
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [8D1-8C7] recognizes Collagen III
Tested Reactivity	Hu
Tested Application	ELISA, ICC/IF, IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	8D1-8C7
Isotype	IgG1, kappa
Target Name	Collagen III
Species	Human
Immunogen	Purified type III collagen extracted from Human placenta.
Conjugation	Un-conjugated
Alternate Names	EDS4A; Collagen alpha-1(III) chain

Application Instructions

Application table	Application	Dilution
	ELISA	1:2000
	ICC/IF	1:40
	IHC-P	1:500
	WB	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Purified.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	COL3A1
Gene Full Name	collagen, type III, alpha 1
Background	This gene encodes the pro-alpha1 chains of type III collagen, a fibrillar collagen that is found in extensible connective tissues such as skin, lung, uterus, intestine and the vascular system, frequently in association with type I collagen. Mutations in this gene are associated with Ehlers-Danlos syndrome types IV, and with aortic and arterial aneurysms. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene. [provided by R. Dagleish, Feb 2008]
Function	Collagen type III occurs in most soft connective tissues along with type I collagen. Involved in regulation of cortical development. Is the major ligand of ADGRG1 in the developing brain and binding to ADGRG1 inhibits neuronal migration and activates the RhoA pathway by coupling ADGRG1 to GNA13 and possibly GNA12. [UniProt]
Calculated Mw	139 kDa
PTM	Proline residues at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains. O-linked glycan consists of a Glc-Gal disaccharide bound to the oxygen atom of a post-translationally added hydroxyl group. [UniProt]
Cellular Localization	Secreted, extracellular space, extracellular matrix. [UniProt]

Images



ARG23989 anti-Collagen III antibody [8D1-8C7] IHC-P image

Immunohistochemistry: Paraffin-embedded Human skin tissue stained with ARG23989 anti-Collagen III antibody [8D1-8C7].
