

ARG30342 Neuroendocrine Tumor IHC Antibody Panel

Package: 1 kit
Store at: -20°C

Component


Cat. No.	Component Name	Host clonality	Reactivity	Application	Package
ARG66347	anti-Ki-67 antibody [SQab1897]	Rabbit mAb	Hu	IHC-P	50 µl
ARG66772	anti-Chromogranin A antibody [SQab20205]	Rabbit mAb	Hu	IHC-P	50 µl
ARG66629	anti-Synaptophysin antibody [SQab19147]	Rabbit mAb	Hu, Ms, Rat	IHC-P, WB	50 µl
ARG66820	anti-CD56 / NCAM antibody [SQab20232]	Rabbit mAb	Hu	IHC-P	50 µl

Properties

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note For laboratory research only, not for drug, diagnostic or other use.

Images




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human tonsil tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human esophageal cancer tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human cervical cancer tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human diffuse large B-cell lymphoma stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human esophageal cancer tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human tonsil tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66772 anti-Chromogranin A antibody [SQab20205] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human colon tissue. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). The tissue section was stained with ARG66772 anti-Chromogranin A antibody [SQab20205] at 18°C - 25°C for 30 minutes.




ARG66629 anti-Synaptophysin antibody [SQab19147] WB image

Western blot: 20 µg of Mouse brain and Rat brain lysates stained with ARG66629 anti-Synaptophysin antibody [SQab19147] at 1:500 dilution.



ARG66629 anti-Synaptophysin antibody [SQab19147] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human glioma tissue stained with ARG66629 anti-Synaptophysin antibody [SQab19147]. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0).



ARG66820 anti-CD56 / NCAM antibody [SQab20232] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human glioma tissue. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). The tissue section was stained with ARG66820 anti-CD56 / NCAM antibody [SQab20232].