

**ARG40169**  
anti-eIF3E antibodyPackage: 50 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes eIF3E
Tested Reactivity	Hu
Predict Reactivity	Cow, Dog, Pig
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	eIF3E
Species	Human
Immunogen	Synthetic peptide around the C-terminus of Human eIF3E. (C-KLNQNSRSEAPN, NP_001559.1)
Conjugation	Un-conjugated
Alternate Names	Viral integration site protein INT-6 homolog; EIF3-P48; EIF3S6; eIF3e; Eukaryotic translation initiation factor 3 subunit 6; eIF3-p46; INT6; Eukaryotic translation initiation factor 3 subunit E; eIF-3 p48

### Application Instructions

Application table	Application	Dilution
	WB	0.03 - 0.1 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	50 kDa	

### Properties

Form	Liquid
Purification	Ammonium sulphate precipitation followed by affinity purification with immunogen.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note**

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


---

Gene Symbol	EIF3E
Gene Full Name	eukaryotic translation initiation factor 3, subunit E
Function	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA <sup>i</sup> and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins. [UniProt]
Calculated Mw	52 kDa
Cellular Localization	Cytoplasm. Nucleus, PML body. [UniProt]

## Images

---

Daudi



### ARG40169 anti-eIF3E antibody WB image

Western blot: 35 µg of Daudi cell lysate (in RIPA buffer) stained with ARG40169 anti-eIF3E antibody at 0.03 µg/ml dilution and incubated at RT for 1 hour.