

ARG40280 anti-NQO1 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes NQO1
Tested Reactivity	Hu, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	NQO1
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 242-274 of Human NQO1. (EVQDEEKNNKFGLSVGHHLGKSIPTDNQIKARK)
Conjugation	Un-conjugated
Alternate Names	DTD; QR1; DHQU; DIA4; NMOR1; NMORI; NAD(P)H dehydrogenase [quinone] 1; EC 1.6.5.2; Azoreductase; DT-diaphorase; DTD; Menadione reductase; NAD(P)H:quinone oxidoreductase 1; Phylloquinone reductase; Quinone reductase 1; QR1

Application Instructions

Application table	Application	Dilution
	WB	0.1 - 0.5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 30 kDa	

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na ₂ HPO ₄ , 0.9% NaCl, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	NQO1
Gene Full Name	NAD(P)H dehydrogenase, quinone 1
Background	This gene is a member of the NAD(P)H dehydrogenase (quinone) family and encodes a cytoplasmic 2-electron reductase. This FAD-binding protein forms homodimers and reduces quinones to hydroquinones. This protein's enzymatic activity prevents the one electron reduction of quinones that results in the production of radical species. Mutations in this gene have been associated with tardive dyskinesia (TD), an increased risk of hematotoxicity after exposure to benzene, and susceptibility to various forms of cancer. Altered expression of this protein has been seen in many tumors and is also associated with Alzheimer's disease (AD). Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]
Function	The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinons involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis. [UniProt]
Highlight	Related products: NQO1 antibodies ; NQO1 Duos / Panels ; Anti-Rabbit IgG secondary antibodies ; Related news: Keap1-Nrf2-ARE antibody panel is launched
Calculated Mw	31 kDa
Cellular Localization	Cytoplasm. [UniProt]

Images



ARG40280 anti-NQO1 antibody WB image

Western blot: 50 µg of Rat liver, 50 µg of Rat lung, 40 µg of HeLa, 40 µg of A549, 40 µg of MDA-MB-231, 40 µg of SW620 and 40 µg of 22Rv1 whole cell lysates stained with ARG40280 anti-NQO1 antibody at 0.5 µg/ml dilution.