

## ARG40477 anti-MMP16 antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes MMP16
Tested Reactivity	Hu, Rat
Predict Reactivity	Ms, Hm
Tested Application	IHC-Fr, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MMP16
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 582-598 of Human MMP16. (YTVFQFKRKGTPRHILY)
Conjugation	Un-conjugated
Alternate Names	MT-MMP 3; MT-MMP2; MT3-MMP; MTMMP3; MT-MMP3; C8orf57; EC 3.4.24.-; MMP-X2; Membrane-type matrix metalloproteinase 3; Membrane-type-3 matrix metalloproteinase; MMP-16; MT3MMP; Matrix metalloproteinase-16

### Application Instructions

Application table	Application	Dilution
	IHC-Fr	1:200 - 1:1000
	IHC-P	0.5 - 1 µg/ml
	WB	0.1 - 0.5 µg/ml

**Application Note**  
IHC-P: Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min.  
\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.9% NaCl, 0.05% Thimerosal, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Thimerosal and 0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated


freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note** For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	MMP16
Gene Full Name	matrix metalloproteinase 16 (membrane-inserted)
Background	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The encoded protein activates MMP2 by cleavage. This gene was once referred to as MT-MMP2, but was renamed as MT-MMP3 or MMP16. [provided by RefSeq, Oct 2010]
Function	Endopeptidase that degrades various components of the extracellular matrix, such as collagen type III and fibronectin. Activates progelatinase A. Involved in the matrix remodeling of blood vessels. Isoform short cleaves fibronectin and also collagen type III, but at lower rate. It has no effect on type I, II, IV and V collagen. However, upon interaction with CSPG4, it may be involved in degradation and invasion of type I collagen by melanoma cells. [UniProt]
Calculated Mw	70 kDa
PTM	The precursor is cleaved by a furin endopeptidase. [UniProt]
Cellular Localization	Isoform Long: Cell membrane; Single-pass type I membrane protein; Extracellular side. Note=Localized at the cell surface of melanoma cells. Isoform Short: Secreted, extracellular space, extracellular matrix. Cell surface. Note=Localized at the cell surface of melanoma cells. [UniProt]

## Images




ARG40477 anti-MMP16 antibody IHC-Fr image

Immunohistochemistry: Frozen section of Human placenta tissue. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG40477 anti-MMP16 antibody at 1 µg/ml dilution, overnight at 4°C.




ARG40477 anti-MMP16 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human rectal cancer tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG40477 anti-MMP16 antibody at 1 µg/ml dilution, overnight at 4°C.




#### ARG40477 anti-MMP16 antibody WB image

Western blot: 50  $\mu$ g of samples under reducing conditions. HeLa and COLO320 whole cell lysates stained with ARG40477 anti-MMP16 antibody at 0.5  $\mu$ g/ml, overnight at 4°C.



#### ARG40477 anti-MMP16 antibody WB image

Western blot: 30  $\mu$ g of sample under reducing conditions. Human heart lysate stained with ARG40477 anti-MMP16 antibody at 0.5  $\mu$ g/ml dilution, overnight at 4°C.



#### ARG40477 anti-MMP16 antibody IHC-Fr image

Immunohistochemistry: Frozen section of Rat intestine tissue. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG40477 anti-MMP16 antibody at 1  $\mu$ g/ml dilution, overnight at 4°C.