

ARG41377
anti-Phosphotyrosine antibodyPackage: 100 µl
Store at: -20°C

Summary

| | |
|---------------------|---|
| Product Description | Rabbit Polyclonal antibody recognizes Phosphotyrosine |
| Tested Reactivity | Hu, Ms, Rat |
| Tested Application | Dot, FACS, ICC/IF, IP, WB |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | Phosphotyrosine |
| Species | Others |
| Immunogen | Phosphotyrosine. |
| Conjugation | Un-conjugated |


Application Instructions

| Application table | Application | Dilution |
|-------------------|-------------|-----------------|
| | Dot | Assay-dependent |
| | FACS | 1:50 |
| | ICC/IF | 1:50 - 1:200 |
| | IP | 1:50 |
| | WB | 1:500 - 1:2000 |

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

| | |
|---------------------|---|
| Form | Liquid |
| Purification | Affinity purified. |
| Buffer | PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol. |
| Preservative | 0.02% Sodium azide |
| Stabilizer | 50% Glycerol |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |



ARG41377 anti-Phosphotyrosine antibody ICC/IF image

Immunofluorescence: A431 cells treated with pervanadate. Cells were stained with ARG41377 anti-Phosphotyrosine antibody.



ARG41377 anti-Phosphotyrosine antibody WB image

Western blot: Jurkat cell lysate stained with ARG41377 anti-Phosphotyrosine antibody.