

ARG41442 anti-Cyclin E2 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Cyclin E2
Tested Reactivity	Hu, Ms
Tested Application	FACS, ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Cyclin E2
Species	Human
Immunogen	Synthetic peptide of Human Cyclin E2.
Conjugation	Un-conjugated
Alternate Names	CYCE2; G1/S-specific cyclin-E2

Application Instructions

Application table	Application	Dilution
	FACS	1:20
	ICC/IF	1:100 - 1:500
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	
Observed Size	~ 50 kDa	

Properties


Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	CCNE2
Gene Full Name	cyclin E2
Background	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2. This cyclin has been shown to specifically interact with CIP/KIP family of CDK inhibitors, and plays a role in cell cycle G1/S transition. The expression of this gene peaks at the G1-S phase and exhibits a pattern of tissue specificity distinct from that of cyclin E1. A significantly increased expression level of this gene was observed in tumor-derived cells. [provided by RefSeq, Jul 2008]
Function	Essential for the control of the cell cycle at the late G1 and early S phase. [UniProt]
Calculated Mw	47 kDa
PTM	Phosphorylation by CDK2 triggers its release from CDK2 and degradation via the ubiquitin proteasome pathway. [UniProt]
Cellular Localization	Nucleus. [UniProt]

Images




ARG41442 anti-Cyclin E2 antibody ICC/IF image

Immunofluorescence: HeLa cells stained with ARG41442 anti-Cyclin E2 antibody.



ARG41442 anti-Cyclin E2 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human breast carcinoma tissue stained with ARG41442 anti-Cyclin E2 antibody.



ARG41442 anti-Cyclin E2 antibody WB image

Western blot: HeLa cell lysate stained with ARG41442 anti-Cyclin E2 antibody.