

ARG41516 anti-OPRD1 / Delta Opioid Receptor antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes OPRD1 / Delta Opioid Receptor
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	OPRD1 / Delta Opioid Receptor
Species	Human
Immunogen	Synthetic peptide of Human OPRD1 / Delta Opioid Receptor.
Conjugation	Un-conjugated
Alternate Names	DOR-1; D-OR-1; Delta-type opioid receptor; OPRD

Application Instructions

Application table	Application	Dilution
	FACS	1:50
	WB	1:1000 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	SH-SY5Y	
Observed Size	~ 42 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	OPRD1
Gene Full Name	opioid receptor, delta 1
Function	G-protein coupled receptor that functions as receptor for endogenous enkephalins and for a subset of other opioids. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling leads to the inhibition of adenylate cyclase activity. Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Plays a role in the perception of pain and in opiate-mediated analgesia. Plays a role in developing analgesic tolerance to morphine. [UniProt]
Calculated Mw	40 kDa
PTM	N-glycosylated. Ubiquitinated. A basal ubiquitination seems not to be related to degradation. Ubiquitination is increased upon formation of OPRM1:OPRD1 oligomers leading to proteasomal degradation; the ubiquitination is diminished by RTP4. [UniProt]
Cellular Localization	Cell membrane; Multi-pass membrane protein. [UniProt]

Images



ARG41516 anti-OPRD1 / Delta Opioid Receptor antibody WB image

Western blot: SH-SY5Y cell lysate stained with ARG41516 anti-OPRD1 / Delta Opioid Receptor antibody.