

ARG41546 anti-CD82 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CD82
Tested Reactivity	Hu, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD82
Species	Human
Immunogen	Synthetic peptide of Human CD82.
Conjugation	Un-conjugated
Alternate Names	Tetraspanin-27; C33 antigen; R2; Tspan-27; TSPAN27; KAI1; CD antigen CD82; CD82 antigen; ST6; 4F9; Suppressor of tumorigenicity 6 protein; SAR2; IA4; Metastasis suppressor Kangai-1; Inducible membrane protein R2; GR15; C33

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Jurkat	
Observed Size	~ 40 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CD82
Gene Full Name	CD82 molecule
Background	This metastasis suppressor gene product is a membrane glycoprotein that is a member of the transmembrane 4 superfamily. Expression of this gene has been shown to be downregulated in tumor progression of human cancers and can be activated by p53 through a consensus binding sequence in the promoter. Its expression and that of p53 are strongly correlated, and the loss of expression of these two proteins is associated with poor survival for prostate cancer patients. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Function	Associates with CD4 or CD8 and delivers costimulatory signals for the TCR/CD3 pathway. [UniProt]
Calculated Mw	30 kDa
Cellular Localization	Membrane; Multi-pass membrane protein. [UniProt]

Images



ARG41546 anti-CD82 antibody WB image

Western blot: Jurkat cell lysate stained with ARG41546 anti-CD82 antibody.