

ARG41597
anti-FSTL3 antibodyPackage: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes FSTL3
Tested Reactivity	Hu, Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	FSTL3
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 27-263 of Human FSTL3 (NP_005851.1).
Conjugation	Un-conjugated
Alternate Names	Follistatin-like protein 3; Follistatin-related gene protein; FLRG; Follistatin-related protein 3; FSRP

Application Instructions

Application table	Application	Dilution
	WB	1:1000 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	SKOV3	
Observed Size	~ 30 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	FSTL3
Gene Full Name	follistatin-like 3 (secreted glycoprotein)
Background	Follistatin-like 3 is a secreted glycoprotein of the follistatin-module-protein family. It may have a role in leukemogenesis. [provided by RefSeq, Jul 2008]
Function	Isoform 1 or the secreted form is a binding and antagonizing protein for members of the TGF-beta family, such as activin, BMP2 and MSTN. Inhibits activin A-, activin B-, BMP2- and MSDT-induced cellular signaling; more effective on activin A than on activin B. Involved in bone formation; inhibits osteoclast differentiation. Involved in hematopoiesis; involved in differentiation of hemopoietic progenitor cells, increases hematopoietic cell adhesion to fibronectin and seems to contribute to the adhesion of hematopoietic precursor cells to the bone marrow stroma. Isoform 2 or the nuclear form is probably involved in transcriptional regulation via interaction with MLLT10. [UniProt]
Calculated Mw	28 kDa
Cellular Localization	Isoform 1: Secreted. Isoform 2: Nucleus. Note=Although alternative initiation has been demonstrated and resulted in different localization, the major source of nuclear FSTL3 appears not to depend on translation initiation at Met-27 according to. [UniProt]

Images



ARG41597 anti-FSTL3 antibody WB image

Western blot: 25 µg of SKOV3 cell lysate stained with ARG41597 anti-FSTL3 antibody at 1:1000 dilution.