

ARG41769 anti-CD71 / Transferrin Receptor antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CD71 / Transferrin Receptor
Tested Reactivity	Hu
Tested Application	FACS, ICC/IF, IHC-Fr, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD71 / Transferrin Receptor
Species	Human
Immunogen	Recombinant protein corresponding to M1-N198 of Human CD71 / Transferrin Receptor.
Conjugation	Un-conjugated
Alternate Names	TFR1; CD antigen CD71; CD71; T9; p90; TR; Trfr; Transferrin receptor protein 1; TRFR; sTfR; TfR1; TfR; TFR

Application Instructions

Application table	Application	Dilution
	FACS	1:150 - 1:500
	ICC/IF	1:200 - 1:1000
	IHC-Fr	1:200 - 1:1000
	IHC-P	1:200 - 1:1000
	WB	1:500 - 1:2000
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0, epitope retrieval solution) for 20 min. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 95 kDa	

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na ₂ HPO ₄ , 0.9% NaCl, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA

Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	TFRC
Gene Full Name	transferrin receptor
Background	This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeq, Sep 2015]
Function	Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. [UniProt]
Calculated Mw	85 kDa
PTM	N- and O-glycosylated, phosphorylated and palmitoylated. The serum form is only glycosylated. Proteolytically cleaved on Arg-100 to produce the soluble serum form (sTfR). Palmitoylated on both Cys-62 and Cys-67. Cys-62 seems to be the major site of palmitoylation. [UniProt]
Cellular Localization	Cell membrane; Single-pass type II membrane protein. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Transferrin receptor protein 1, serum form: Secreted. [UniProt]

Images




ARG41769 anti-CD71 / Transferrin Receptor antibody ICC/IF image

Immunofluorescence: A431 cells were blocked with 10% goat serum and then stained with ARG41769 anti-CD71 / Transferrin Receptor antibody (red) at 2 µg/ml dilution, overnight at 4°C. DAPI (blue) for nuclear staining.




ARG41769 anti-CD71 / Transferrin Receptor antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human placenta tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0, epitope retrieval solution) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG41769 anti-CD71 / Transferrin Receptor antibody at 1 $\mu\text{g}/\text{ml}$ dilution, overnight at 4°C.




ARG41769 anti-CD71 / Transferrin Receptor antibody WB image

Western blot: 50 μg of samples under reducing conditions. HeLa, Jurkat, Raji, HL-60, K562, HepG2, Human placenta and CEM whole cell lysates stained with ARG41769 anti-CD71 / Transferrin Receptor antibody at 0.5 $\mu\text{g}/\text{ml}$ dilution, overnight at 4°C.



ARG41769 anti-CD71 / Transferrin Receptor antibody FACS image


Flow Cytometry: SiHa cells were blocked with 10% normal goat serum and then stained with ARG41769 anti-CD71 / Transferrin Receptor antibody (blue) at 1 $\mu\text{g}/10^6$ cells for 30 min at 20°C, followed by incubation with DyLight®488 labelled secondary antibody. Isotype control antibody (green) was Rabbit IgG (1 $\mu\text{g}/10^6$ cells) used under the same conditions. Unlabelled sample (red) was also used as a control.



ARG41769 anti-CD71 / Transferrin Receptor antibody IHC-Fr image

Immunohistochemistry: Frozen section of Human placenta tissue. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG41769 anti-CD71 / Transferrin Receptor antibody at 1 $\mu\text{g}/\text{ml}$ dilution, overnight at 4°C.

ARG41769 anti-CD71 / Transferrin Receptor antibody FACS image



Flow Cytometry: U87 cells were blocked with 10% normal goat serum and then stained with ARG41769 anti-CD71 / Transferrin Receptor antibody (blue) at $1 \mu\text{g}/10^6$ cells for 30 min at 20°C , followed by incubation with DyLight[®]488 labelled secondary antibody. Isotype control antibody (green) was Rabbit IgG ($1 \mu\text{g}/10^6$ cells) used under the same conditions. Unlabelled sample (red) was also used as a control.