

ARG41851 anti-FBXO32 antibody [M314]

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rat Monoclonal antibody [M314] recognizes FBXO32
Tested Reactivity	Hu
Tested Application	WB
Host	Rat
Clonality	Monoclonal
Clone	M314
Isotype	IgG
Target Name	FBXO32
Species	Mouse
Immunogen	Synthetic peptide (coupled to carrier protein) corresponding to aa. 23-35 of Mouse FBXO32.
Conjugation	Un-conjugated
Alternate Names	Muscle atrophy F-box protein; Atrogin-1; F-box only protein 32; Fbx32; MAFbx

Application Instructions

Application table	Application	Dilution
	WB	1:500

Application Note
WB: Antibody is suggested to be diluted in 5% skimmed milk/Tris buffer with 0.04% Tween20 and incubated for 1 hour at room temperature.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.05% Sodium azide, 50% Glycerol and 1 mg/ml BSA.
Preservative	0.05% Sodium azide
Stabilizer	50% Glycerol and 1 mg/ml BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	FBXO32
Gene Full Name	F-box protein 32
Background	This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class and contains an F-box domain. This protein is highly expressed during muscle atrophy, whereas mice deficient in this gene were found to be resistant to atrophy. This protein is thus a potential drug target for the treatment of muscle atrophy. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jun 2011]
Function	Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Probably recognizes and binds to phosphorylated target proteins during skeletal muscle atrophy. Recognizes TERF1. [UniProt]
Calculated Mw	42 kDa
Cellular Localization	Cytoplasm. Nucleus. Note=Shuttles between cytoplasm and the nucleus. [UniProt]

Images



ARG41851 anti-FBXO32 antibody [M314] WB image

Western blot: Human FBXO32 recombinant protein stained with ARG41851 anti-FBXO32 antibody [M314] at 1:1000, 1:2000 and 1:4000 dilution.