

**ARG42030**  
anti-Sall4 antibodyPackage: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes Sall4
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Sall4
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 853-1053 of Human Sall4 (NP_065169.1).
Conjugation	Un-conjugated
Alternate Names	dJ1112F19.1; Zinc finger protein 797; DRRS; Zinc finger protein SALL4; Sal-like protein 4; HSAL4; ZNF797

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HepG2	
Observed Size	~ 75 kDa	

### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note**

For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

---

Gene Symbol	SALL4
Gene Full Name	spalt-like transcription factor 4
Background	The protein encoded by this gene may be a zinc finger transcription factor. Defects in this gene are a cause of Duane-radial ray syndrome (DRRS). [provided by RefSeq, Jul 2008]
Function	Transcription factor with a key role in the maintenance and self-renewal of embryonic and hematopoietic stem cells. [UniProt]
Calculated Mw	112 kDa
PTM	Isoform SALL4B exists primarily as a ubiquitinated form.  Sumoylation with both SUMO1 and SUMO2 regulates the stability, subcellular localization, transcriptional activity, and may reduce interaction with POU5F1/OCT4. [UniProt]
Cellular Localization	Cytoplasm. Nucleus. [UniProt]

## Images

---



HepG2

ARG42030 anti-Sall4 antibody WB image

Western blot: 25 µg of HepG2 cell lysate stained with ARG42030 anti-Sall4 antibody at 1:1000 dilution.