

ARG42033 anti-EGLN1 + EGLN2 antibody

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes EGLN1 + EGLN2
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	EGLN1 + EGLN2
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-426 of Human EGLN1 (NP_071334.1).
Conjugation	Un-conjugated
Alternate Names	EGLN1: HPH-2; HIF-PH2; Egl nine homolog 1; HIF-prolyl hydroxylase 2; HPH2; ZMYND6; C1orf12; HALAH; PHD2; Prolyl hydroxylase domain-containing protein 2; Hypoxia-inducible factor prolyl hydroxylase 2; SM-20; HIFPH2; EC 1.14.11.29; SM20; ECT3 EGLN2: EIT6; PHD1; EIT-6; HPH-1; HPH-3; HIFPH1; HIF-PH1

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Rat heart	
Observed Size	~ 46 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	EGLN1; EGLN2
Gene Full Name	egl-9 family hypoxia-inducible factor 1 egl-9 family hypoxia inducible factor 2
Background	<p>EGLN1: The protein encoded by this gene catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. HIF is a transcriptional complex that plays a central role in mammalian oxygen homeostasis. This protein functions as a cellular oxygen sensor, and under normal oxygen concentration, modification by prolyl hydroxylation is a key regulatory event that targets HIF subunits for proteasomal destruction via the von Hippel-Lindau ubiquitylation complex. Mutations in this gene are associated with erythrocytosis familial type 3 (ECYT3). [provided by RefSeq, Nov 2009]</p> <p>EGLN2: The hypoxia inducible factor (HIF) is a transcriptional complex that is involved in oxygen homeostasis. At normal oxygen levels, the alpha subunit of HIF is targeted for degradation by prolyl hydroxylation. This gene encodes an enzyme responsible for this post-translational modification. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the upstream RAB4B (RAB4B, member RAS oncogene family) gene. [provided by RefSeq, Feb 2011]</p>
Function	<p>EGLN1: Cellular oxygen sensor that catalyzes, under normoxic conditions, the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates a specific proline found in each of the oxygen-dependent degradation (ODD) domains (N-terminal, NODD, and C-terminal, CODD) of HIF1A. Also hydroxylates HIF2A. Has a preference for the CODD site for both HIF1A and HIF1B. Hydroxylated HIFs are then targeted for proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Under hypoxic conditions, the hydroxylation reaction is attenuated allowing HIFs to escape degradation resulting in their translocation to the nucleus, heterodimerization with HIF1B, and increased expression of hypoxia-inducible genes. EGLN1 is the most important isozyme under normoxia and, through regulating the stability of HIF1, involved in various hypoxia-influenced processes such as angiogenesis in retinal and cardiac functionality. Target proteins are preferentially recognized via a LXXLAP motif. [UniProt]</p> <p>EGLN2: Cellular oxygen sensor that catalyzes, under normoxic conditions, the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates a specific proline found in each of the oxygen-dependent degradation (ODD) domains (N-terminal, NODD, and C-terminal, CODD) of HIF1A. Also hydroxylates HIF2A. Has a preference for the CODD site for both HIF1A and HIF2A. Hydroxylated HIFs are then targeted for proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Under hypoxic conditions, the hydroxylation reaction is attenuated allowing HIFs to escape degradation resulting in their translocation to the nucleus, heterodimerization with HIF1B, and increased expression of hypoxia-inducible genes. EGLN2 is involved in regulating hypoxia tolerance and apoptosis in cardiac and skeletal muscle. Also regulates susceptibility to normoxic oxidative neuronal death. Links oxygen sensing to cell cycle and primary cilia formation by hydroxylating the critical centrosome component CEP192 which promotes its ubiquitination and subsequent proteasomal degradation. Hydroxylates IKBKB, mediating NF-kappaB activation in hypoxic conditions. Target proteins are preferentially recognized via a LXXLAP motif.</p>
Calculated Mw	46 kDa
PTM	EGLN1: S-nitrosylation inhibits the enzyme activity up to 60% under aerobic conditions. Chelation of Fe(2+) has no effect on the S-nitrosylation. It is uncertain whether nitrosylation occurs on Cys-323 or Cys-326. [UniProt]
Cellular Localization	<p>EGLN1: Cytoplasm. Nucleus. Note=Mainly cytoplasmic. Shuttles between the nucleus and cytoplasm (PubMed:19631610). Nuclear export requires functional XPO1. [UniProt]</p> <p>EGLN2: Nucleus. [UniProt]</p>




ARG42033 anti-EGLN1 + EGLN2 antibody ICC/IF image

Immunofluorescence: L929 cells stained with ARG42033 anti-EGLN1 + EGLN2 antibody at 1:100 dilution. DAPI (blue) for nuclear staining.



ARG42033 anti-EGLN1 + EGLN2 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human colon tissue stained with ARG42033 anti-EGLN1 + EGLN2 antibody at 1:100 dilution.



ARG42033 anti-EGLN1 + EGLN2 antibody WB image

Western blot: 25 µg of Rat heart lysate stained with ARG42033 anti-EGLN1 + EGLN2 antibody at 1:3000 dilution.
