

ARG42350 anti-CD274 / PD-L1 antibody [29E.2A3] (low endotoxin)

Package: 100 µg
Store at: -20°C

Summary

Product Description	Azide free and low endotoxin Mouse Monoclonal antibody [29E.2A3] recognizes CD274 / PD-L1
Tested Reactivity	Hu, NHuPrm
Tested Application	FACS, FuncSt, IHC-Fr
Specificity	The mouse monoclonal antibody 29E.2A3 recognizes an extracellular epitope of CD274 / PD-L1 (also known as B7-H1), a 40 kDa type I transmembrane protein expressed by dendritic cells, activated T cells, activated monocytes, and in various tissues, above all in heart and skeletal muscle, placenta and lung, and in many cancer cells, including T cell lymphomas, melanomas, and glioblastomas.
Host	Mouse
Clonality	Monoclonal
Clone	29E.2A3
Isotype	IgG2b, kappa
Target Name	CD274 / PD-L1
Species	Human
Immunogen	Full length Human CD274.
Conjugation	Un-conjugated
Alternate Names	Programmed cell death 1 ligand 1; B7-H1; B7H1; PDL1; PDCD1 ligand 1; B7 homolog 1; PD-L1; CD antigen CD274; PDCD1L1; B7-H; Programmed death ligand 1; PDCD1LG1

Application Instructions

Application table	Application	Dilution
	FACS	1 - 4 µg/ml
	FuncSt	Assay-dependent
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purification with Protein A.
Purification Note	0.2 µm filter sterilized. Endotoxin level is less than 0.01 EU/µg of the protein.
Buffer	PBS
Concentration	1 mg/ml

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	CD274
Gene Full Name	CD274 molecule
Background	This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]
Function	<p>Plays a critical role in induction and maintenance of immune tolerance to self (PubMed:11015443, PubMed:28813417, PubMed:28813410). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed:11015443, PubMed:28813417, PubMed:28813410). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed:10581077).</p> <p>The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed:28813417, PubMed:28813410). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (By similarity). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (By similarity). [UniProt]</p>
Highlight	<p>Related products: PD-L1 antibodies; PD-L1 ELISA Kits; Anti-Mouse IgG secondary antibodies;</p> <p>Related news: Examining CTL/NK-mediated cytotoxicity by ELISA</p>
Calculated Mw	33 kDa
Cellular Localization	Cell membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Recycling endosome membrane; Single-pass type I membrane protein. Note=Associates with CMTM6 at recycling endosomes, where it is protected from being targeted for lysosomal degradation. Isoform 1: Cell membrane; Single-pass type I membrane protein. Isoform 2: Endomembrane system; Single-pass type I membrane protein. [UniProt]



ARG42350 anti-CD274 / PD-L1 antibody [29E.2A3] (low endotoxin)
FACS image

Flow Cytometry: Human PHA stimulated peripheral blood mononuclear cell suspension stained with ARG42350 anti-CD274 / PD-L1 antibody [29E.2A3] (low endotoxin) at 4 µg/ml dilution, followed by APC-conjugated Goat anti-Mouse antibody.



ARG42350 anti-CD274 / PD-L1 antibody [29E.2A3] (low endotoxin)
FACS image

Flow Cytometry: Separation of PHA stimulated peripheral blood mononuclear cell suspension (red-filled) from unstimulated cells (black-dashed). Cells were stained with ARG42350 anti-CD274 / PD-L1 antibody [29E.2A3] (low endotoxin) at 4 µg/ml dilution, followed by APC-conjugated Goat anti-Mouse antibody.