

ARG42389
anti-KIR2DL1 + KIR2DS5 + KIR2DS1 + KIRDS3 antibody [HP-MA4]Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [HP-MA4] recognizes KIR2DL1 + KIR2DS5 + KIR2DS1 + KIRDS3
Tested Reactivity	Hu
Tested Application	FACS, IP
Specificity	The mouse monoclonal antibody HP-MA4 recognizes an extracellular epitope of CD158 isoforms KIR2DL1 (CD158a), KIR2DS5 (CD158g), KIR2DS1 (CD158h), and KIRDS3. It does not recognize the isoforms CD158b1,d,f,i,j.
Host	Mouse
Clonality	Monoclonal
Clone	HP-MA4
Isotype	IgG2b, kappa
Target Name	KIR2DL1 + KIR2DS5 + KIR2DS1 + KIRDS3
Species	Human
Immunogen	Human NK cell line LB2.
Conjugation	Un-conjugated
Alternate Names	p58.1 MHC class-I-specific NK receptor; NKAT1; Killer cell immunoglobulin-like receptor 2DL1; NKAT-1; p58 natural killer cell receptor clones CL-42/47.11; CD158A; KIR-K64; CD158 antigen-like family member A; p58.1; NKAT; p58 NK receptor CL-42/47.11; MHC class I NK cell receptor; Natural killer-associated transcript 1; CD antigen CD158a; KIR221

Application Instructions

Application table	Application	Dilution
	FACS	1 - 4 µg/ml
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS and 15 mM Sodium azide.
Preservative	15 mM Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot


and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	KIR2DL1
Gene Full Name	killer cell immunoglobulin-like receptor, two domains, long cytoplasmic tail, 1
Background	Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008]
Function	Receptor on natural killer (NK) cells for some HLA-C alleles such as w4 and w6. Inhibits the activity of NK cells thus preventing cell lysis. [UniProt]
Calculated Mw	39 kDa
Cellular Localization	Cell membrane; Single-pass type I membrane protein. [UniProt]

Images



ARG42389 anti-KIR2DL1 + KIR2DS5 + KIR2DS1 + KIRDS3 antibody [HP-MA4] FACS image

Flow Cytometry: Human peripheral blood stained with ARG42389 anti-KIR2DL1 + KIR2DS5 + KIR2DS1 + KIRDS3 antibody [HP-MA4], followed by APC-conjugated Goat anti-Mouse antibody. Samples were co-stained with anti-CD3 antibody (PE).