

ARG42888 anti-RPS6KA6 / RSK4 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes RPS6KA6 / RSK4
Tested Reactivity	Hu
Tested Application	FACS, ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	lgG
Target Name	RPS6KA6 / RSK4
Species	Human
Immunogen	Synthetic peptide derived from Human RPS6KA6 / RSK4.
Conjugation	Un-conjugated
Alternate Names	Ribosomal S6 kinase 4; pp90RSK4; p90-RSK 6; p90RSK6; RSK-4; PP90RSK4; S6K-alpha-6; EC 2.7.11.1; 90 kDa ribosomal protein S6 kinase 6; RSK4; Ribosomal protein S6 kinase alpha-6

Application Instructions

Application table	Application	Dilution
	FACS	1:50
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	SH-SY5Y	
Observed Size	~ 88 kDa	

Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	RPS6KA6
Gene Full Name	ribosomal protein S6 kinase, 90kDa, polypeptide 6
Background	This gene encodes a member of ribosomal S6 kinase family, serine-threonine protein kinases which are regulated by growth factors. The encoded protein may be distinct from other members of this family, however, as studies suggest it is not growth factor dependent and may not participate in the same signaling pathways. [provided by RefSeq, Jan 2010]
Function	Constitutively active serine/threonine-protein kinase that exhibits growth-factor-independent kinase activity and that may participate in p53/TP53-dependent cell growth arrest signaling and play an inhibitory role during embryogenesis. [UniProt]
Calculated Mw	84 kDa
РТМ	Phosphorylated at Ser-232, Ser-372, and Ser-389 in serum-starved cells. [UniProt]
Cellular Localization	Cytoplasm, cytosol. Nucleus. Note=Predominantly cytosolic. [UniProt]

Images

