

Product datasheet

info@arigobio.com

ARG42900 anti-SFRS3 / SRSF3 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes SFRS3 / SRSF3

Tested Reactivity Hu, Ms, Rat

Tested Application ICC/IF, IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name SFRS3 / SRSF3

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 1-90 of Human SFRS3 / SRSF3 (NP_003008.1).

Conjugation Un-conjugated

Alternate Names Splicing factor, arginine/serine-rich 3; Serine/arginine-rich splicing factor 3; SFRS3; Pre-mRNA-splicing

factor SRP20; SRp20

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	U-87MG	
Observed Size	~ 22 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol SRSF3

Gene Full Name serine/arginine-rich splicing factor 3

Background The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-mRNA

splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two transcript variants, one protein-coding and the

other non-coding, have been found for this gene. [provided by RefSeq, Sep 2010]

Function Splicing factor that specifically promotes exon-inclusion during alternative splicing (PubMed:26876937).

Interaction with YTHDC1, a RNA-binding protein that recognizes and binds N6-methyladenosine (m6A)-containing RNAs, promotes recruitment of SRSF3 to its mRNA-binding elements adjacent to m6A sites, leading to exon-inclusion during alternative splicing (PubMed:26876937). Also functions as export adapter involved in mRNA nuclear export (PubMed:11336712, PubMed:18364396, PubMed:28984244). Binds mRNA which is thought to be transferred to the NXF1-NXT1 heterodimer for export (TAP/NXF1 pathway); enhances NXF1-NXT1 RNA-binding activity (PubMed:11336712, PubMed:18364396). Involved in nuclear export of m6A-containing mRNAs via interaction with YTHDC1 facilitates m6A-containing mRNA-binding to both SRSF3 and NXF1, promoting mRNA nuclear export (PubMed:28984244). RNA-binding is semi-sequence specific (PubMed:17036044). [UniProt]

Calculated Mw 19 kDa


PTM Phosphorylated by CLK1, CLK2, CLK3 and CLK4. Extensively phosphorylated on serine residues in the RS

domain. [UniProt]

Cellular Localization Nucleus. Nucleus speckle. Cytoplasm. Note=Recruited to nuclear speckles following interaction with


YTHDC1. [UniProt]

Images




ARG42900 anti-SFRS3 / SRSF3 antibody ICC/IF image

Immunofluorescence: NIH/3T3 cells stained with ARG42900 anti-SFRS3 / SRSF3 antibody at 1:100 dilution.



ARG42900 anti-SFRS3 / SRSF3 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human mammary cancer tissue stained with ARG42900 anti-SFRS3 / SRSF3 antibody at 1:100 dilution.



ARG42900 anti-SFRS3 / SRSF3 antibody WB image

Western blot: 25 μg of U-87MG cell lysate stained with ARG42900 anti-SFRS3 / SRSF3 antibody at 1:3000 dilution.