

# Product datasheet

info@arigobio.com

# ARG42913 anti-UCHL5 / UCH37 antibody

Package: 100 μl Store at: -20°C

### Summary

Product Description Mouse Monoclonal antibody recognizes UCHL5 / UCH37

Tested Reactivity Hu
Tested Application WB

Host Mouse

Clonality Monoclonal
Clone 854CT5.5.6

Isotype IgG1

Target Name UCHL5 / UCH37

Species Human

Immunogen Purified His-tagged Human UCHL5 / UCH37 protein.

Conjugation Un-conjugated

Alternate Names UCH-L5; CGI-70; UCH37; INO80R; Ubiquitin thioesterase L5; Ubiquitin C-terminal hydrolase UCH37;

Ubiquitin carboxyl-terminal hydrolase isozyme L5; EC 3.4.19.12

#### **Application Instructions**

Application table	Application	Dilution
	WB	1:2000 - 1:4000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	
Observed Size	~ 35 kDa	

#### **Properties**

Form Liquid

Purification Purification with Protein G.

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol UCHL5

Gene Full Name ubiquitin carboxyl-terminal hydrolase L5

Function Protease that specifically cleaves 'Lys-48'-linked polyubiquitin chains. Deubiquitinating enzyme

associated with the 19S regulatory subunit of the 26S proteasome. Putative regulatory component of the INO80 complex; however is inactive in the INO80 complex and is activated by a transient interaction


of the INO80 complex with the proteasome via ADRM1. [UniProt]

Calculated Mw 38 kDa

Cellular Localization Cytoplasm. Nucleus. Note=Associates with the proteasome 19S subunit in the cytoplasm. Associates

with the INO80 complex in the nucleus. [UniProt]

## **Images**



Western blot: 20  $\mu g$  of HeLa whole cell lysate stained with ARG42913 anti-UCHL5 / UCH37 antibody at 1:2000 - 1:4000 dilution.

