

Product datasheet

info@arigobio.com

ARG43257 anti-PI3 Kinase p85 alpha antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes PI3 Kinase p85 alpha

Tested Reactivity Hu, Rat, Hm

Predict Reactivity Ms

Tested Application FACS, ICC/IF, IP, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name PI3 Kinase p85 alpha

Species Human

Immunogen Synthetic peptide of Human PI3 Kinase p85 alpha.

Conjugation Un-conjugated

Alternate Names GRB1; PI3-kinase subunit p85-alpha; Phosphatidylinositol 3-kinase regulatory subunit alpha; IMD36;

Ptdlns-3-kinase regulatory subunit alpha; p85-ALPHA; p85; AGM7; Ptdlns-3-kinase regulatory subunit p85-alpha; Pl3-kinase regulatory subunit alpha; Pl3-kinase r

3-kinase 85 kDa regulatory subunit alpha

Application Instructions

FACS 1:20 ICC/IF 1:50 IP 1:20 WB 1:1000 Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	Application table	Application	Dilution
IP 1:20 WB 1:1000 Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations		FACS	1:20
WB 1:1000 Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations		ICC/IF	1:50
Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations		IP	1:20
		WB	1:1000
should be determined by the scientist.	Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control Rat brain	Positive Control	Rat brain	
Observed Size ~ 86 kDa	Observed Size	~ 86 kDa	

Properties

Form	Liquid	
Purification	Affinity purified.	
Buffer	50 mM Tris-Glycine (pH 7.4), 150 mM NaCl, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.	
Preservative	0.01% Sodium azide	

Stabilizer 40% Glycerol and 0.05% BSA

Concentration Batch dependent

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

> and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol PIK3R1

Gene Full Name phosphoinositide-3-kinase, regulatory subunit 1 (alpha)

Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime Background

> position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit. Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in four transcript variants encoding different

isoforms. [provided by RefSeq, Jun 2011]

Function Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an

> adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed:17626883, PubMed:19805105, PubMed:7518429). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the

liver and hence plays a role in glucose tolerance improvement (PubMed:20348923). [UniProt]

Calculated Mw 84 kDa


PTM Polyubiquitinated in T-cells by CBLB; which does not promote proteasomal degradation but impairs

association with CD28 and CD3Z upon T-cell activation.

Phosphorylated. Tyrosine phosphorylated in response to signaling by FGFR1, FGFR2, FGFR3 and FGFR4. Phosphorylated by CSF1R. Phosphorylated by ERBB4. Phosphorylated on tyrosine residues by TEK/TIE2. Dephosphorylated by PTPRJ. Phosphorylated by PIK3CA at Ser-608; phosphorylation is stimulated by insulin and PDGF. The relevance of phosphorylation by PIK3CA is however unclear (By similarity).

Phosphorylated in response to KIT and KITLG/SCF. Phosphorylated by FGR. [UniProt]

Images



ARG43257 anti-PI3 Kinase p85 alpha antibody WB image

Western blot: Rat brain lysate stained with ARG43257 anti-PI3 Kinase p85 alpha antibody at 1:1000 dilution.