

Product datasheet

info@arigobio.com

ARG43433 anti-GPR56 antibody

Package: $50 \mu l$ Store at: $-20 ^{\circ}C$

Summary

Product Description Mouse Monoclonal antibody recognizes GPR56

Tested Reactivity Hu

Tested Application ICC/IF, WB
Host Mouse

Clonality Monoclonal Isotype IgG1, kappa

Target Name GPR56
Species Human

Immunogen Recombinant protein of Human GPR56.

Conjugation Un-conjugated

Alternate Names BPPR; G-protein coupled receptor 56; C; GPR56 subunit alpha; BFPP; GPR56 NT; Protein TM7XN1;

TM7XN1; GPR56 seven-transmembrane subunit; GPR56 extracellular subunit; N; GPR56 subunit beta;

GPR56 7TM; GPR56; TM7LN4; GPR56 CT

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:25
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	U87-MG	
Observed Size	~ 75 kDa	

Properties

Form Liquid

Purification Purification with Protein G.

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol

ADGRG1

Gene Full Name

adhesion G protein-coupled receptor G1

Background

This gene encodes a member of the G protein-coupled receptor family and regulates brain cortical patterning. The encoded protein binds specifically to transglutaminase 2, a component of tissue and tumor stroma implicated as an inhibitor of tumor progression. Mutations in this gene are associated with a brain malformation known as bilateral frontoparietal polymicrogyria. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]

Function

Receptor involved in cell adhesion and probably in cell-cell interactions. Mediates cell matrix adhesion in developing neurons and hematopoietic stem cells. Receptor for collagen III/COL3A1 in the developing brain and involved in regulation of cortical development, specifically in maintenance of the pial basement membrane integrity and in cortical lamination (By similarity). Binding to the COL3A1 ligand inhibits neuronal migration and activates the RhoA pathway by coupling to GNA13 and possibly GNA12 (PubMed:22238662). Plays a role in the maintenance of hematopoietic stem cells and/or leukemia stem cells in bone marrow niche (By similarity). Plays a critical role in cancer progression by inhibiting VEGFA production threreby inhibiting angiogenesis through a signaling pathway mediated by PRKCA (PubMed:16757564, PubMed:21724588). Plays an essential role in testis development (By similarity).

[ADGRG1 N-terminal fragment]: Plays a critical role in cancer progression by activating VEGFA production and angiogenesis through a signaling pathway mediated by PRKCA (PubMed:21724588). [UniProt]

Calculated Mw

78 kDa

PTM

Autoproteolytically cleaved into 2 fragments; the large extracellular N-terminal fragment (ADGRG1 NT) and the membrane-bound C-terminal fragment (ADGRG1 CT) predominantly remain associated and non-covalently linked. Shedding to yield the secreted ADGRG1 N-terminal fragment seems to involve metalloprotease(s) (PubMed:22333914).


N-glycosylated. Contains sialic acid residues.

Ubiquitinated. Undergoes polyubiquitination upon activation. [UniProt]

Cellular Localization

Cell membrane; Multi-pass membrane protein. ADGRG1 N-terminal fragment: Secreted. ADGRG1 C-terminal fragment: Membrane raft. Note=Interaction with its ligand COL3A1 leads to the release of ADGRG1 NT from the membrane and triggers the association of ADGRG1 CT with lipid rafts. [UniProt]

Images



ARG43433 anti-GPR56 antibody WB image

Western blot: $20~\mu g$ of U87-MG whole cell lysate stained with ARG43433 anti-GPR56 antibody at 1:1000 dilution.