

## Product datasheet

info@arigobio.com

# ARG43479 anti-PXDN / VPO1 antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Product Description Rabbit Polyclonal antibody recognizes PXDN / VPO1.

Tested Reactivity Hu, Ms
Tested Application WB
Host Rabbit

**Clonality** Polyclonal

Isotype IgG

Target Name PXDN / VPO1

Species Human

Immunogen Purified recombinant protein corresponding to human PXDN / VPO1.

Conjugation Un-conjugated

Alternate Names PXN; VPO; MG50; PRG2; COPOA; D2S448; D2S448E

#### **Application Instructions**

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recomm should be determined by the sci	nended starting dilutions and the optimal dilutions or concentrations lentist.

### **Properties**

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

|--|

www.arigobio.com arigo.nuts about antibodies 1/2

Gene Full Name peroxidasin

Background This gene encodes a heme-containing peroxidase that is secreted into the extracellular matrix. It is

involved in extracellular matrix formation, and may function in the physiological and pathological fibrogenic response in fibrotic kidney. Mutations in this gene cause corneal opacification and other ocular anomalies, and also microphthalmia and anterior segment dysgenesis. [provided by RefSeq, Aug

2014]

Function Displays low peroxidase activity and is likely to participate in H(2)O(2) metabolism and peroxidative

reactions in the cardiovascular system. Plays a role in extracellular matrix formation. [UniProt]