

## Product datasheet

info@arigobio.com

# ARG43544 anti-HNF1 beta / vHNF1 antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Product Description Rabbit Polyclonal antibody recognizes HNF1 beta / vHNF1.

Tested Reactivity Hu
Tested Application WB

Host Rabbit

**Clonality** Polyclonal

Isotype IgG

Target Name HNF1 beta / vHNF1

Species Human

Immunogen Synthetic peptide derived from human HNF1 beta / vHNF1

Conjugation Un-conjugated

Alternate Names HNF1; LFB1; TCF1; MODY3; TCF-1; HNF-1A; IDDM20

#### **Application Instructions**

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### **Properties**

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol	HNF1A			
-------------	-------	--	--	--

www.arigobio.com arigo.nuts about antibodies 1/2

Gene Full Name HNF1 homeobox A

Background The protein encoded by this gene is a transcription factor required for the expression of several liver-

specific genes. The encoded protein functions as a homodimer and binds to the inverted palindrome 5'-GTTAATNATTAAC-3'. Defects in this gene are a cause of maturity onset diabetes of the young type 3 (MODY3) and also can result in the appearance of hepatic adenomas. Alternative splicing results in

multiple transcript variants encoding different isoforms. [provided by RefSeq, Apr 2015]

Function Transcriptional activator that regulates the tissue specific expression of multiple genes, especially in

pancreatic islet cells and in liver. Required for the expression of several liver specific genes. Binds to the

inverted palindrome 5'-GTTAATNATTAAC-3'. [UniProt]