

**ARG43767**  
anti-FNDC5 antibodyPackage: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes FNDC5
Tested Reactivity	Hu, Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	FNDC5
Species	Human
Immunogen	Synthetic peptide corresponding to Human FNDC5.
Conjugation	Un-conjugated
Alternate Names	irisin; FRCP2; Fibronectin type III domain-containing protein 5; Fibronectin type III repeat-containing protein 2

### Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	20-25 kDa	


### Properties

Form	Liquid
Purification	Affinity Purified
Buffer	50 mM Tris-Glycine (pH 7.4), 0.15M NaCl, 0.05% BSA, 40% Glycerol and 0.01% Sodium azide.
Preservative	0.01% Sodium azide
Stabilizer	0.05% BSA and 40% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	FNDC5
Gene Full Name	fibronectin type III domain containing 5
Background	This gene encodes a secreted protein that is released from muscle cells during exercise. The encoded protein may participate in the development of brown fat. Translation of the precursor protein initiates at a non-AUG start codon at a position that is conserved as an AUG start codon in other organisms. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2013]
Function	Irisin: Contrary to mouse, may not be involved in the beneficial effects of muscular exercise, nor in the induction of browning of human white adipose tissue. [UniProt]
Calculated Mw	23 kDa
PTM	The extracellular domain is cleaved and released from the cell membrane. N-Glycosylated. [UniProt]
Cellular Localization	Cell membrane. [UniProt]

## Images



ARG43767 anti-FNDC5 antibody WB image

Western blot: HeLa and 3T3 cell lysate stained with ARG43767 anti-FNDC5 antibody at 1:1000 dilution.



ARG43767 anti-FNDC5 antibody WB image

Western blot: C6 cell lysate stained with ARG43767 anti-FNDC5 antibody at 1:800 dilution.