

ARG44153 anti-CD172a + CD172b antibody [SE5A5] (APC)

Package: 100 tests
Store at: 4°C

Summary

Product Description	APC-conjugated Mouse Monoclonal antibody [SE5A5] recognizes CD172a + CD172b
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The mouse monoclonal antibody SE5A5 recognizes a common extracellular epitope on human CD172a and CD172b antigens (approx. 90 kDa and approx. 50 kDa, respectively), although its reactivity with CD172a is higher.
Host	Mouse
Clonality	Monoclonal
Clone	SE5A5
Isotype	IgG1
Target Name	CD172a + CD172b
Species	Human
Immunogen	NIH-3T3 cells / Human CD172a.
Conjugation	APC
Alternate Names	CD172A; p84; SHPS1; SHPS-1; CD172 antigen-like family member A; Sirp-alpha-3; Sirp-alpha-1; BIT; MYD-1; MFR; Bit; PTPNS1; CD antigen CD172a; Inhibitory receptor SHPS-1; SIRP; MyD-1 antigen; Sirp-alpha-2; Tyrosine-protein phosphatase non-receptor type substrate 1; Signal-regulatory protein alpha-1; Signal-regulatory protein alpha-2; Signal-regulatory protein alpha-3; Macrophage fusion receptor; Brain Ig-like molecule with tyrosine-based activation motifs; P84; SHP substrate 1

Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>10 µl / 100 µl of whole blood or 10⁶ cells</td></tr></tbody></table>	Application	Dilution	FACS	10 µl / 100 µl of whole blood or 10 ⁶ cells
Application	Dilution				
FACS	10 µl / 100 µl of whole blood or 10 ⁶ cells				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				

Properties

Form	Liquid
Purification	Purified
Buffer	PBS(pH 7.4) and 15 mM Sodium azide.
Preservative	15 mM Sodium azide
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	SIRPA
Gene Full Name	signal-regulatory protein alpha
Background	<p>The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phospho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phosphatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal transduction mediated by various growth factor receptors. CD47 has been demonstrated to be a ligand for this receptor protein. This gene and its product share very high similarity with several other members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene. [provided by RefSeq, Jul 2008]</p>
Function	<p>Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function (By similarity). Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. [UniProt]</p>
Calculated Mw	55 kDa
PTM	<p>N-glycosylated.</p> <p>Phosphorylated on tyrosine residues in response to stimulation with EGF, growth hormone, insulin and PDGF. Dephosphorylated by PTPN11. [UniProt]</p>
Cellular Localization	Membrane; Single-pass type I membrane protein. [UniProt]

Images



ARG44153 anti-CD172a + CD172b antibody [SE5A5] (APC) FACS image

Flow Cytometry: Human peripheral whole blood stained with ARG44153 anti-CD172a + CD172b antibody [SE5A5] (APC) at 10 μ l / 100 μ l of whole blood dilution.