

Product datasheet

info@arigobio.com

ARG44182 anti-MOCS1 antibody

Package: 50 μg Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes MOCS1

Tested Reactivity Hu, Ms, Rat

Tested Application FACS, ICC/IF, IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name MOCS1
Species Human

Immunogen Recombinant protein of Human MOCS1

Conjugation Un-conjugated

Alternate Names MOCS1; Molybdenum Cofactor Synthesis 1; MOCOD; Molybdenum Cofactor Synthesis-Step 1 Protein A-

B; Molybdenum Cofactor Biosynthesis Protein 1; Cell Migration-Inducing Gene 11 Protein; Molybdenum

Cofactor Biosynthesis Protein A; MOCS1A Enzyme; MOCS1A; MOCS1B; MIG11

Application Instructions

Application table	Application	Dilution
	FACS	1-3 μg/1x10^6 cells
	ICC/IF	5 μg/ml
	IHC-P	2-5 μg/ml
	WB	0.25-0.5 μg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Affinity purification with immunogen.

Buffer 0.9% NaCl, 0.2% Na2HPO4, 0.05% Sodium azide and 5% BSA.

Preservative 0.05% Sodium azide

Stabilizer 5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol MOCS1

Gene Full Name Molybdenum Cofactor Synthesis 1

Background Molybdenum cofactor biosynthesis is a conserved pathway leading to the biological activation of

molybdenum. The protein encoded by this gene is involved in this pathway. This gene was originally thought to produce a bicistronic mRNA with the potential to produce two proteins (MOCS1A and MOCS1B) from adjacent open reading frames. However, only the first open reading frame (MOCS1A) has been found to encode a protein from the putative bicistronic mRNA, whereas additional splice variants are likely to produce a fusion between the two open reading frames. This gene is defective in patients with molybdenum cofactor deficiency, type A. A related pseudogene has been identified on

chromosome 16.

Function Isoform MOCS1A and isoform MOCS1B probably form a complex that catalyzes the conversion of

5'-GTP to cyclic pyranopterin monophosphate (cPMP). MOCS1A catalyzes the cyclization of GTP to (8S)-3',8-cyclo-7,8-dihydroguanosine 5'-triphosphate and MOCS1B catalyzes the subsequent conversion


of (8S)-3',8-cyclo-7,8-dihydroguanosine 5'-triphosphate to cPMP.

Calculated Mw 70 kDa

PTM Acetylation, Phosphoprotein


Cellular Localization Cytosol, Nucleus

Images




ARG44182 anti-MOCS1 antibody IHC-P image

Immunohistochemistry: Human liver cancer stained with ARG44182 anti-MOCS1 antibody at 2 $\mu g/mL$ dilution.




ARG44182 anti-MOCS1 antibody ICC/IF image

Immunofluorescence: HepG2 stained with ARG44182 anti-MOCS1 antibody at 5 $\mu g/mL$ dilution.




ARG44182 anti-MOCS1 antibody WB image

Western blot: HepG2, Jurkat and Caco-2 stained with ARG44182 anti-MOCS1 antibody at 0.5 $\mu g/mL$ dilution.




ARG44182 anti-MOCS1 antibody FACS image

Flow Cytometry: K562 stained with ARG44182 anti-MOCS1 antibody at 1 μ g/1x10^6 cells dilution.




ARG44182 anti-MOCS1 antibody WB image

Western blot: rat liver and PC-12 stained with ARG44182 anti-MOCS1 antibody at 0.5 $\mu g/mL$ dilution.




ARG44182 anti-MOCS1 antibody WB image

Western blot: Mouse liver and RAW264.7 stained with ARG44182 anti-MOCS1 antibody at 0.5 $\mu g/mL$ dilution.



ARG44182 anti-MOCS1 antibody IHC-P image

Immunohistochemistry: Human testicular germ cell tumors stained with ARG44182 anti-MOCS1 antibody at 2 μ g/mL dilution.



ARG44182 anti-MOCS1 antibody IHC-P image

Immunohistochemistry: Human colorectal adenocarcinoma stained with ARG44182 anti-MOCS1 antibody at 2 μ g/mL dilution.