

## ARG44241 anti-Langerin / CD207 antibody [2G3]

Package: 100 µg  
Store at: 4°C

### Summary

Product Description	Mouse Monoclonal antibody [2G3] recognizes Langerin / CD207
Tested Reactivity	Hu, NHuPrm
Tested Application	ELISA, FACS, ICC/IF, IHC-Fr
Host	Mouse
Clonality	Monoclonal
Clone	2G3
Isotype	IgG1, lambda
Target Name	Langerin / CD207
Species	Human
Immunogen	Human CD207 Fusion protein
Conjugation	Un-conjugated
Alternate Names	CD207; CD207 Molecule; CLEC4K; C-Type Lectin Domain Family 4 Member K Langerin; Langerhans Cell Specific C-Type Lectin; CD207 Molecule, Langerin; CD207 Antigen, Langerin; C-Type Lectin Domain Family 4, Member K; CD207 Antigen

### Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	FACS	0.5-4 µg/ml
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS (pH 7.4) and 15 mM Sodium azide.
Preservative	15 mM Sodium azide
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Do not freeze. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation

---

Gene Symbol	CD207
Gene Full Name	CD207 Molecule
Background	The protein encoded by this gene is expressed only in Langerhans cells which are immature dendritic cells of the epidermis and mucosa. It is localized in the Birbeck granules, organelles present in the cytoplasm of Langerhans cells and consisting of superimposed and zippered membranes. It is a C-type lectin with mannose binding specificity, and it has been proposed that mannose binding by this protein leads to internalization of antigen into Birbeck granules and providing access to a nonclassical antigen-processing pathway. Mutations in this gene result in Birbeck granules deficiency or loss of sugar binding activity.
Function	Calcium-dependent lectin displaying mannose-binding specificity. Induces the formation of Birbeck granules (BGs); is a potent regulator of membrane superimposition and zippering. Binds to sulfated as well as mannosylated glycans, keratan sulfate (KS) and beta-glucans. Facilitates uptake of antigens and is involved in the routing and/or processing of antigen for presentation to T cells. Major receptor on primary Langerhans cells for <i>Candida</i> species, <i>Saccharomyces</i> species, and <i>Malassezia furfur</i> . Protects against human immunodeficiency virus-1 (HIV-1) infection. Binds to high-mannose structures present on the envelope glycoprotein which is followed by subsequent targeting of the virus to the Birbeck granules leading to its rapid degradation.
PTM	Disulfide bond, Glycoprotein
Cellular Localization	Membrane

## Images



ARG44241 anti-Langerin / CD207 antibody [2G3] FACS image

Flow Cytometry: Human whole blood stained with ARG44241 anti-Langerin / CD207 antibody [2G3] at 0.56 µg/ml dilution.