

Product datasheet

info@arigobio.com

ARG44313 anti-CSRNP1 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes CSRNP1

Tested Reactivity Hu

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name CSRNP1
Species Human

ImmunogenSynthetic peptideConjugationUn-conjugated

Alternate Names CSRNP1; Cysteine And Serine Rich Nuclear Protein 1; TAIP-3; FAM130B; URAX1; AXUD1; Cysteine/Serine-

Rich Nuclear Protein 1; TGF-Beta-Induced Apoptosis Protein 3; Axin-1 Up-Regulated Gene 1 Protein;

AXIN1 Up-Regulated 1; CSRNP-1

Application Instructions

Application table	Application	Dilution
	WB	1:500-1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Antigen Affinity Purified

Buffer PBS with 0.02% Sodium azide

Preservative 0.02% Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol CSRNP1

Gene Full Name Cysteine And Serine Rich Nuclear Protein 1

Background This gene encodes a protein that localizes to the nucleus and expression of this gene is induced in

response to elevated levels of axin. The Wnt signalling pathway, which is negatively regulated by axin, is important in axis formation in early development and impaired regulation of this signalling pathway is often involved in tumors. A decreased level of expression of this gene in tumors compared to the level of expression in their corresponding normal tissues suggests that this gene product has a tumor

suppressor function. Alternative splicing results in multiple transcript variants.

Function Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity.

Calculated Mw 64 kDa

Cellular Localization Nucleus