

**ARG44629**  
anti-BIRC2 / cIAP1 antibodyPackage: 50 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody recognizes BIRC2 / cIAP1
Tested Reactivity	Hu
Tested Application	IP
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Target Name	BIRC2 / cIAP1
Species	Human
Conjugation	Un-conjugated
Alternate Names	BIRC2; Baculoviral IAP Repeat Containing 2; C-IAP1; RNF48; MIHB; Hiap-2; CIAP1; API1; Baculoviral IAP Repeat-Containing Protein 2; RING-Type E3 Ubiquitin Transferase BIRC2; TNFR2-TRAF-Signaling Complex Protein 2; NFR2-TRAF Signalling Complex Protein; Cellular Inhibitor Of Apoptosis 1; Inhibitor Of Apoptosis Protein 2; RING Finger Protein 48; Apoptosis Inhibitor 1; IAP Homolog B; HIAP2; Baculoviral IAP Repeat-Containing 2; EC 2.3.2.27; HIAP-2; IAP-2

### Application Instructions

Application table	Application	Dilution
	IP	10 µg/mL

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


### Properties

Form	Liquid
Purification	Protein A purification
Buffer	PBS with 0.09% sodium azide
Preservative	0.09% sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	BIRC2
Gene Full Name	Baculoviral IAP Repeat Containing
Background	The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. This encoded protein inhibits apoptosis induced by serum deprivation and menadione, a potent inducer of free radicals. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012]
Function	Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, mitogenic kinase signaling, and cell proliferation, as well as cell invasion and metastasis. Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and regulates both canonical and non-canonical NF-kappa-B signaling by acting in opposite directions: acts as a positive regulator of the canonical pathway and suppresses constitutive activation of non-canonical NF-kappa-B signaling. The target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, RIPK2, RIPK3, RIPK4, CASP3, CASP7, CASP8, TRAF2, DIABLO/SMAC, MAP3K14/NIK, MAP3K5/ASK1, IKBKG/NEMO, IKBKE and MXD1/MAD1. Can also function as an E3 ubiquitin-protein ligase of the NEDD8 conjugation pathway, targeting effector caspases for neddylation and inactivation. Acts as an important regulator of innate immune signaling via regulation of Toll-like receptors (TLRs), Nodlike receptors (NLRs) and RIG-I like receptors (RLRs), collectively referred to as pattern recognition receptors (PRRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase-independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8. Can stimulate the transcriptional activity of E2F1. Plays a role in the modulation of the cell cycle.
PTM	Ubl conjugation. [UniProt]
Cellular Localization	Cytoplasm; Nucleus. [UniProt]

## Images



ARG44629 anti-BIRC2 / cIAP1 antibody IP image

Immunoprecipitation: HEK293 lysate immunoprecipitated with 2.5  $\mu$ g of ARG44629 anti-BIRC2 / cIAP1 antibody.