

ARG45014 anti-IgA2 antibody [RM125]

Package: 50 µg
Store at: -20°C

Summary

| | |
|---------------------|---|
| Product Description | Rabbit Monoclonal antibody [RM125] recognizes IgA2. |
| Tested Reactivity | Hu |
| Tested Application | ELISA, ICC/IF, IHC-P |
| Specificity | This antibody reacts to human IgA2. No cross reactivity with human IgA1, IgG, IgM, IgD, or IgE. |
| Host | Rabbit |
| Clonality | Monoclonal |
| Clone | RM125 |
| Isotype | IgG |
| Target Name | IgA2 |
| Species | Human |
| Immunogen | Human IgA2 |
| Target Ig | IgA2 |
| Conjugation | Un-conjugated |

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|---|
| | ELISA | 50 - 200 ng/well (for Capture); 0.05 - 0.2 µg/mL (for Detection) |
| | ICC/IF | 0.5 - 2 µg/mL |
| | IHC-P | 0.1 - 1 µg/ml |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

| | |
|---------------------|--|
| Buffer | PBS with 50% Glycerol, 1% BSA and 0.09% sodium azide |
| Preservative | 0.09% sodium azide |
| Stabilizer | 50% Glycerol, 1% BSA and 0.09% |
| Concentration | 1 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

Note

For laboratory research only, not for drug, diagnostic or other use.