

ARG52296 anti-GABAA Receptor beta 2 antibody

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes GABAA Receptor beta 2
Tested Reactivity	Rat
Predict Reactivity	Hu, Ms, Dog, NHuPrm
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	GABAA Receptor beta 2
Species	Rat
Immunogen	Fusion protein from the cytoplasmic loop of the beta 2 subunit
Conjugation	Un-conjugated
Alternate Names	A; Gamma-aminobutyric acid receptor subunit beta-2; GABA

Application Instructions

Application table	Application	Dilution
	WB	1:1,000

Application Note
Specific for the ~55k β2-subunit of the GABAA receptor in Western blots.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties


Form	Liquid
Purification	Affinity Purified
Buffer	10 mM HEPES (pH 7.5), 150 mM NaCl, 0.1 mg/ml BSA and 50% Glycerol
Stabilizer	0.1 mg/ml BSA, 50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links [GeneID: 25451 Rat](#)

Gene Symbol	GABRB2
Gene Full Name	gamma-aminobutyric acid (GABA) A receptor, beta 2
Background	Gamma-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl ⁻ channel associated with the GABAA receptor (GABAA-R) subtype. GABAA-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABAA-R is a multimeric subunit complex. To date six α s, four β s and four γ s, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for α - and β -subunits results in the expression of functional GABAA-Rs sensitive to GABA. However, coexpression of a γ -subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different α - subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2004; Pörtl et al., 2003).
Research Area	Neuroscience antibody
Calculated Mw	59 kDa

Images



ARG52296 anti-GABAA Receptor beta 2 antibody WB image

Western Blot: 5-7 ug of rat cerebellum showing specific immunolabeling of the ~55k beta 2-subunit of the GABAA-R stained with GABAA Receptor beta 2 antibody (ARG52296).