

## ARG52346 anti-Neurofilament NF-H phospho (KSP site) antibody [NAP4]

Package: 50 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [NAP4] recognizes Neurofilament NF-H when KSP site was phosphorylated
Tested Reactivity	Ms, Rat, Cow, Pig
Predict Reactivity	Hu
Tested Application	IHC-P, WB
Specificity	NAP4 recognizes phosphorylated NF-H KSP sequences but not non-phosphorylated KSP sequences.
Host	Mouse
Clonality	Monoclonal
Clone	NAP4
Isotype	IgG1
Target Name	Neurofilament NF-H
Species	Bovine
Immunogen	Purified bovine NF-H
Conjugation	Un-conjugated
Alternate Names	Neurofilament heavy polypeptide; 200 kDa neurofilament protein; NF-H; Neurofilament triplet H protein; NFH

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:500 - 1:1000
	WB	1:5000 - 1:10000

Application Note	<p>Specific for the ~200k Neurofilament H protein. It recognizes phosphorylated NF-H KSP (lysine-serine-proline) type sequences. In some species there is some cross-reactivity with the related phosphorylated KSP sequences found in the related neurofilament subunit NF-M. It recognizes neurofilaments in frozen sections in tissue culture and in formalin fixed sections.</p> <p>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.</p>
------------------	--

### Properties

Form	Liquid
Purification	Total IgG fraction
Buffer	Total IgG fraction and 10 mM Sodium azide
Preservative	10 mM Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot


and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note** For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Database links	<a href="#">GeneID: 100156492 Pig</a> <a href="#">GeneID: 100625407 Pig</a> <a href="#">GeneID: 380684 Mouse</a> <a href="#">Swiss-port # P19246 Mouse</a>
Background	Neurofilaments are the 10nm or intermediate filament proteins found specifically in neurons, and are composed predominantly of three major proteins called NF-L, NF-M and NF-H . NF-H is the neurofilament high or heavy molecular weight polypeptide and runs on SDS-PAGE gels at 200-220 kDa, with some variability across species boundaries. Antibodies to NF-H are useful for identifying neuronal cells and their processes in tissue sections and in tissue culture. NF-H antibodies can also be useful to visualize neurofilament accumulations seen in many neurological diseases, such as Amyotrophic Lateral Sclerosis (Lou Gehrig's disease) (2) and Alzheimer's disease .
Research Area	Neuroscience antibody; Signaling Transduction antibody; Neurofilament antibody; Intermediate Neurofilament antibody
Calculated Mw	112 kDa
PTM	There are a number of repeats of the tripeptide K-S-P, NFH is phosphorylated on a number of the serines in this motif. It is thought that phosphorylation of NFH results in the formation of interfilament cross bridges that are important in the maintenance of axonal caliber. Phosphorylation seems to play a major role in the functioning of the larger neurofilament polypeptides (NF-M and NF-H), the levels of phosphorylation being altered developmentally and coincidentally with a change in the neurofilament function. Phosphorylated in the head and rod regions by the PKC kinase PKN1, leading to the inhibition of polymerization.

## Images




ARG52346 anti-Neurofilament NF-H antibody [NAP4] WB image

Western Blot: Rat cortex lysate showing specific immunolabeling of the ~200k NF-H protein stained with ARG52346 anti-Neurofilament NF-H antibody [NAP4].



#### ARG52346 anti-Neurofilament NF-H antibody [NAP4] IHC image

Immunohistochemistry: Human cerebellar cortex stained with ARG52346 anti-Neurofilament NF-H antibody [NAP4] showing labeling of NF-H (brown) in basket cell axons surrounding the large Purkinje neurons.



#### ARG52346 anti-Neurofilament NF-H antibody [NAP4] WB image

Western blot: Rat brain, Rat spinal cord, Mouse brain, Mouse spinal cord, Pig spinal cord and Cow spinal cord lysates stained with ARG52346 anti-Neurofilament NF-H antibody [NAP4] (green) at 1:10000 dilution.

Strong band at about 200-220 kDa corresponds to the major phosphorylated form of the NF-H subunit. A minor band at about 160 kDa is the non-phosphorylated NF-H. Smaller proteolytic fragments of NF-H are also detected in spinal cord preparations.