

ARG52463 anti-Tyrosine Hydroxylase phospho (Ser31) antibody

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Tyrosine Hydroxylase phospho (Ser31)
Tested Reactivity	Ms, Rat
Predict Reactivity	NHuPrm
Tested Application	ICC/IF, IHC, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Tyrosine Hydroxylase
Species	Rat
Immunogen	Synthetic phospho-peptide corresponding to amino acid residues surrounding Ser31 conjugated to KLH
Conjugation	Un-conjugated
Alternate Names	DYT14; TYH; Tyrosine 3-monoxygenase; Tyrosine 3-hydroxylase; TH; DYT5b; EC 1.14.16.2

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:1000
	IHC	frozen sections: 1:1000
	WB	1:1000
Application Note	Specific for the ~60k tyrosine hydroxylase protein phosphorylated at Ser31. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity Purified
Buffer	10 mM HEPES (pH 7.5), 150 mM NaCl, 0.1 mg/ml BSA and 50% Glycerol
Stabilizer	0.1 mg/ml BSA, 50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 21823 Mouse GeneID: 25085 Rat Swiss-port # P04177 Rat Swiss-port # P24529 Mouse
Gene Symbol	TH
Gene Full Name	tyrosine hydroxylase
Background	<p>Tyrosine hydroxylase (TH) is the rate-limiting enzyme in the synthesis of the catecholamines dopamine and norepinephrine. TH antibodies can therefore be used as markers for dopaminergic and noradrenergic neurons in a variety of applications including depression, schizophrenia, Parkinson's disease and drug abuse (Kish et al., 2001; Zhu et al., 2000; Zhu et al., 1999). TH antibodies can also be used to explore basic mechanisms of dopamine and norepinephrine signaling (Witkovsky et al., 2000; Salvatore et al., 2001; Dunkley et al., 2004). The activity of TH is also regulated by phosphorylation (Haycock et al., 1982; Haycock et al., 1992; Jedynak et al., 2002). Phospho-specific antibodies for the phosphorylation sites on TH can be used to great effect in studying this regulation and in identifying the cells in which TH phosphorylation occurs.</p>
Highlight	<p>Related Antibody Duos and Panels: ARG30212 Phospho Tyrosine Hydroxylase Antibody Panel (Total, pS31, pS40) Related products: Tyrosine Hydroxylase antibodies: Tyrosine Hydroxylase Duos / Panels: Anti-Rabbit IgG secondary antibodies: Related news: Astrocyte-to-neuron conversion for Parkinson's disease treatment</p>
Research Area	Cancer antibody; Metabolism antibody; Neuroscience antibody
Calculated Mw	59 kDa

Images



ARG52463 anti-Tyrosine Hydroxylase phospho (Ser31) antibody WB image

Western blot: PC-12 cells incubated in the absence (Control) and presence of okadaic acid (OA, 1 μ M for 60 min) showing specific immunolabeling of the \sim 60 kDa TH phosphorylated at Ser31 stained with ARG52463 anti-Tyrosine Hydroxylase phospho (Ser31) antibody.