

**ARG53739**  
anti-CD99 antibody [12E7]Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [12E7] recognizes CD99
Tested Reactivity	Hu
Tested Application	FACS, ICC/IF, IHC-P, IP
Host	Mouse
Clonality	Monoclonal
Clone	12E7
Isotype	IgG1
Target Name	CD99
Immunogen	Acute lymphocytic leukemia T-cells.
Conjugation	Un-conjugated
Alternate Names	12E7; CD99 antigen; MIC2X; MIC2Y; CD antigen CD99; MSK5X; Protein MIC2; MIC2; T-cell surface glycoprotein E2; HBA71; E2 antigen

### Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	ICC/IF	Assay-dependent
	IHC-P	Assay-dependent
	IP	Assay-dependent

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification	Purified Antibody
Buffer	1X PBS and 0.1% Sodium azide
Preservative	0.1% Sodium azide
Concentration	0.2 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 4267 Human</a> <a href="#">Swiss-port # P14209 Human</a>
Gene Symbol	CD99
Gene Full Name	CD99 molecule
Background	CD99 is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. There is a related pseudogene located immediately adjacent to this locus. [provided by RefSeq, Mar 2016]
Function	CD99 involved in T-cell adhesion processes and in spontaneous rosette formation with erythrocytes. Plays a role in a late step of leukocyte extravasation helping leukocytes to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1. Involved in T-cell adhesion processes. [UniProt]
Research Area	Cancer antibody; Immune System antibody; Signaling Transduction antibody
Calculated Mw	19 kDa
PTM	Extensively O-glycosylated.