

ARG53818 anti-CD3 epsilon antibody [MEM-57] (APC)

Package: 100 tests, 50 tests
Store at: 4°C

Summary

Product Description	APC-conjugated Mouse Monoclonal antibody [MEM-57] recognizes CD3 epsilon
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The clone MEM-57 reacts with gamma-epsilon and delta-epsilon dimers of human CD3 complex, a part of a bigger multisubunit T cell receptor complex (CD3/TCR) expressed on peripheral blood T lymphocytes and mature thymocytes. HLDA IV.; WS Code T 96
Host	Mouse
Clonality	Monoclonal
Clone	MEM-57
Isotype	IgG2a
Target Name	CD3 epsilon (activation epitope)
Species	Human
Immunogen	Human thymocytes and T lymphocytes.
Conjugation	APC
Alternate Names	CD3E; CD3 Epsilon Subunit Of T-Cell Receptor Complex; T-Cell Surface Glycoprotein CD3 Epsilon Chain; CD3e Antigen, Epsilon Polypeptide (TiT3 Complex); T-Cell Surface Antigen T3/Leu-4 Epsilon Chain; CD3e Molecule, Epsilon (CD3-TCR Complex); CD3-Epsilon; CD3epsilon

Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>10 µl / 10⁶ cells</td></tr></tbody></table>	Application	Dilution	FACS	10 µl / 10 ⁶ cells
Application	Dilution				
FACS	10 µl / 10 ⁶ cells				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				

Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Preservative	15 mM Sodium azide
Stabilizer	0.2% (w/v) high-grade protease free BSA
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid


repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 915 Human Swiss-port # P04234 Human
Gene Symbol	CD3E
Gene Full Name	CD3 Epsilon Subunit Of T-Cell Receptor Complex
Background	The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also been linked to a susceptibility to type I diabetes in women.
Function	Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways.
Calculated Mw	19 kDa
Cellular Localization	Cell membrane, Membrane

Images



ARG53818 anti-CD3 epsilon antibody [MEM-57] (APC) FACS image

Flow Cytometry: Human peripheral whole blood stained with ARG53818 anti-CD3 epsilon antibody [MEM-57] (APC) (10 µl reagent / 100 µl of peripheral whole blood).



ARG53818 anti-CD3 epsilon antibody [MEM-57] (APC) FACS image

Flow Cytometry: Separation of human CD3 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed). Human peripheral whole blood stained with ARG53818 anti-CD3 epsilon antibody [MEM-57] (APC) (10 μ l reagent / 100 μ l of peripheral whole blood).