

ARG53897 anti-CD72 antibody [3F3] (PE)

Package: 100 tests
Store at: 4°C

Summary

Product Description	PE-conjugated Mouse Monoclonal antibody [3F3] recognizes CD72
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The clone 3F3 reacts with CD72, a 39-43 kDa type II membrane glycoprotein (C-type lectin family). CD72 is a pan-B cell marker expressed throughout the B lymphocytes differentiation with the exception of plasma cells; it is also present on follicular dendritic cells. HLDA V; WS Code B CD72.5 HLDA VI; WS Code B CD72.1 HLDA VI; WS Code 6 BP 84
Host	Mouse
Clonality	Monoclonal
Clone	3F3
Isotype	IgG2b
Target Name	CD72
Species	Human
Immunogen	Normal human lymphocytes from a lymph node.
Conjugation	PE
Alternate Names	CD antigen CD72; LYB2; B-cell differentiation antigen CD72; Lyb-2; CD72b

Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>20 µl / 10⁶ cells</td></tr></tbody></table>	Application	Dilution	FACS	20 µl / 10 ⁶ cells
Application	Dilution				
FACS	20 µl / 10 ⁶ cells				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				

Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Preservative	15 mM Sodium azide
Stabilizer	0.2% (w/v) high-grade protease free BSA
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid


repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 971 Human Swiss-port # P21854 Human
Gene Symbol	CD72
Gene Full Name	CD72 molecule
Background	CD72 is a transmembrane glycoprotein expressed as a homodimer especially in B cells, but also in other antigen presenting cells such as dendritic cells and macrophages. Through one of its immunoreceptor tyrosine-based inhibitory motives (ITIMs), CD72 interacts with tyrosine phosphatase SHP-1, thereby suppressing B cell responsiveness. Binding of CD72 with its ligand CD100 (Sema4D) prevents BCR association and phosphorylation of CD72 and results in dissociation of SHP-1 from CD72, thus enables B cell activation.
Function	Plays a role in B-cell proliferation and differentiation. [UniProt]
Research Area	Developmental Biology antibody; Immune System antibody
Calculated Mw	40 kDa
PTM	Phosphorylated upon engagement of the B-cell receptor, probably by LYN or SYK. Phosphorylation at Tyr-7 is important for interaction with PTPN6/SHP-1 (By similarity).

Images



ARG53897 anti-CD72 antibody [3F3] (PE) FACS image

Flow Cytometry: Human peripheral blood cells stained with ARG53897 anti-CD72 antibody [3F3] (PE).