

Product datasheet

info@arigobio.com

ARG54888 anti-Slug antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes Slug

Tested Reactivity Hu

Predict Reactivity Ms, Rat, Bov

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name Slug

Species Human

Immunogen KLH-conjugated synthetic peptide corresponding to aa. 1-30 (N-terminus) of Human Slug.

Conjugation Un-conjugated

Alternate Names Protein snail homolog 2; SNAIL2; WS2D; Neural crest transcription factor Slug; Zinc finger protein

SNAI2; SLUG; SLUGH1

Application Instructions

Application table	Application	Dilution	
	WB	1:1000	
Application Note		* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A2058		

Properties

Form Liquid

Purification This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis

against PBS.

Buffer PBS and 0.09% (W/V) Sodium azide

Preservative 0.09% (W/V) Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 6591 Human

Swiss-port # O43623 Human

Gene Symbol SNAI2

Gene Full Name snail family zinc finger 2

Background This gene encodes a member of the Snail family of C2H2-type zinc finger transcription factors. The

encoded protein acts as a transcriptional repressor that binds to E-box motifs and is also likely to repress E-cadherin transcription in breast carcinoma. This protein is involved in epithelial-mesenchymal transitions and has antiapoptotic activity. Mutations in this gene may be associated with sporatic cases

of neural tube defects. [provided by RefSeq, Jul 2008]

Function Transcriptional repressor that modulates both activator-dependent and basal transcription. Involved in

the generation and migration of neural crest cells. Plays a role in mediating RAF1-induced

transcriptional repression of the TJ protein, occludin (OCLN) and subsequent oncogenic transformation of epithelial cells (By similarity). Represses BRCA2 expression by binding to its E2-box-containing silencer and recruiting CTBP1 and HDAC1 in breast cells. In epidermal keratinocytes, binds to the E-box in ITGA3 promoter and represses its transcription. Involved in the regulation of ITGB1 and ITGB4 expression and cell adhesion and proliferation in epidermal keratinocytes. Binds to E-box2 domain of BSG and activates its expression during TGFB1-induced epithelial-mesenchymal transition (EMT) in hepatocytes. Represses E-Cadherin/CDH1 transcription via E-box elements. Involved in osteoblast maturation. Binds to RUNX2 and SOC9 promoters and may act as a positive and negative transcription regulator, respectively, in osteoblasts. Binds to CXCL12 promoter via E-box regions in mesenchymal stem cells and osteoblasts. Plays an essential role in TWIST1-induced EMT and its ability to promote

invasion and metastasis. [UniProt]

Research Area Cancer antibody; Controls and Markers antibody; Developmental Biology antibody; Gene Regulation

antibody; Neuroscience antibody

Calculated Mw 30 kDa


PTM GSK3B-mediated phosphorylation results in cytoplasmic localization and degradation.

Cellular Localization Nucleus. Cytoplasm. Note=Observed in discrete foci in interphase nuclei. These nuclear foci do not

overlap with the nucleoli, the SP100 and the HP1 heterochromatin or the coiled body, suggesting SNAI2

is associated with active transcription or active splicing regions

Images



ARG54888 anti-Slug antibody WB image

Western blot: 35 μg of A2058 cell lysate stained with ARG54888 anti-Slug antibody.