

ARG55184 anti-IL17 / IL17A antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes IL17 / IL17A
Tested Reactivity	Hu, Ms
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	IL17 / IL17A
Species	Human
Immunogen	Synthetic peptide (16 aa) within aa. 40-90 of Human IL-17A.
Conjugation	Un-conjugated
Alternate Names	IL-17A; Interleukin-17A; IL-17; CTLA8; IL17; CTLA-8; Cytotoxic T-lymphocyte-associated antigen 8

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	WB	1 - 2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	THP-1 Cell Lysate	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.02% Sodium azide
Preservative	0.02% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 16171 Mouse GeneID: 3605 Human Swiss-port # Q16552 Human Swiss-port # Q62386 Mouse
Gene Symbol	IL17A
Gene Full Name	interleukin 17A
Background	Interleukin 17 (IL-17) is a family of pro-inflammatory cytokines produced by activated T cells and is thought to have a major role in the initiation and perpetuation of rheumatoid arthritis. IL-17 regulates the activities of NF- κ B and mitogen-activated protein kinases such as ERK and JNK. In addition, IL-17 stimulates the expression of IL-6 and cyclooxygenase-2 and enhances the production of nitric oxide. IL-17-producing T helper cells (TH-17 cells) have been the subject of much attention due to the importance of IL-17 in the pathogenesis of autoimmune inflammation. Because of its role in autoimmune diseases, it is thought that targeting the production and action of IL-17 would be beneficial therapeutically in these diseases.
Function	Induces stromal cells to produce proinflammatory and hematopoietic cytokines. Enhances the surface expression of ICAM1/intracellular adhesion molecule 1 in fibroblasts. [UniProt]
Highlight	Related products: IL17 antibodies ; IL17 ELISA Kits ; IL17 recombinant proteins ; Anti-Rabbit IgG secondary antibodies ; Related news: HMGB1, a biomarker and therapeutic target in COVID-19
Research Area	Cancer antibody; Immune System antibody
Calculated Mw	18 kDa
PTM	Found both in glycosylated and nonglycosylated forms.

Images



ARG55184 anti-IL17 / IL17A antibody WB image

Western blot: THP-1 cell lysate stained with ARG55184 anti-IL17 / IL17A antibody at (A) 1 and (B) 2 μ g/ml dilution.