

# Product datasheet

info@arigobio.com

# ARG55899 anti-Hexokinase 2 antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Product Description Mouse Monoclonal antibody recognizes Hexokinase 2

Tested Reactivity Ms, Rat, Mk

Tested Application WB

Host Mouse

**Clonality** Monoclonal

Isotype IgG1

Target Name Hexokinase 2

Species Human

Immunogen Recombinant Human Hexokinase II protein.

Conjugation Un-conjugated

Alternate Names Muscle form hexokinase; HK II; Hexokinase-2; Hexokinase type II; EC 2.7.1.1; HXK2; HKII

## **Application Instructions**

Application table	Application	Dilution
	WB	1:500
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

## **Properties**

Form Liquid

**Purification** Affinity purification with immunogen.

Buffer PBS (pH 7.4), 0.03% Proclin-300 and 50% Glycerol

Preservative 0.03% Proclin-300

Stabilizer 50% Glycerol
Concentration 0.7 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Database links GeneID: 15277 Mouse

GeneID: 25059 Rat

Swiss-port # O08528 Mouse

Swiss-port # P27881 Rat

Gene Symbol HK2

Gene Full Name hexokinase 2

Background Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose

metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-


 $responsive, and studies in \ rat \ suggest \ that \ it \ is \ involved \ in \ the \ increased \ rate \ of \ glycolysis \ seen \ in \ rapidly$ 

growing cancer cells. [provided by RefSeq, Apr 2009]

Calculated Mw 102 kDa

Cellular Localization Mitochondrion outer membrane

## **Images**



## ARG55899 anti-Hexokinase 2 antibody WB image

Western blot: C6 cell lysate stained with ARG55899 anti-Hexokinase 2 antibody at 1:500 dilution.