

ARG56058 anti-Mucin 5AC antibody [1-13M1]

Package: 50 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [1-13M1] recognizes Mucin 5AC
Tested Reactivity	Hu, Ms, Rat
Species Does Not React With	Pig
Tested Application	ELISA
Host	Mouse
Clonality	Monoclonal
Clone	1-13M1
Isotype	IgG1, kappa
Target Name	Mucin 5AC
Species	Human
Immunogen	Mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient.
Conjugation	Un-conjugated
Alternate Names	IeB; Major airway glycoprotein; Lewis B blood group antigen; Tracheobronchial mucin; mucin; Mucin-5AC; LeB; MUC5; TBM; Gastric mucin; MUC-5AC; Mucin-5 subtype AC, tracheobronchial

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS (pH 7.4), 0.05% Sodium azide and 0.1 mg/ml BSA
Preservative	0.05% Sodium azide
Stabilizer	0.1 mg/ml BSA
Concentration	0.2 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 4586 Human
Gene Symbol	MUC5AC
Gene Full Name	mucin 5AC, oligomeric mucus/gel-forming
Function	Gel-forming glycoprotein of gastric and respiratory tract epithelia that protects the mucosa from infection and chemical damage by binding to inhaled microorganisms and particles that are subsequently removed by the mucociliary system. [UniProt]
Calculated Mw	586 kDa
PTM	C-, O- and N-glycosylated. O-glycosylated on the Thr-/Ser-rich tandem repeats. C-mannosylation in the Cys-rich subdomains may be required for proper folding of these regions and for export from the endoplasmic reticulum during biosynthesis. Proteolytic cleavage in the C-terminal is initiated early in the secretory pathway and does not involve a serine protease. The extent of cleavage is increased in the acidic parts of the secretory pathway. Cleavage generates a reactive group which could link the protein to a primary amide.
Cellular Localization	Cytoplasmic