

ARG56931 anti-ISG15 antibody [3E5]

Package: 50 µl
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [3E5] recognizes ISG15
Tested Reactivity	Hu
Tested Application	FACS, WB
Host	Mouse
Clonality	Monoclonal
Clone	3E5
Isotype	IgG1, kappa
Target Name	ISG15
Species	Human
Immunogen	Recombinant fragment around aa. 1-157 of Human ISG15.
Conjugation	Un-conjugated
Alternate Names	G1P2; IP17; UCRP; Interferon-induced 15 kDa protein; Interferon-induced 17 kDa protein; hUCRP; IFI15; Ubiquitin cross-reactive protein; Ubiquitin-like protein ISG15; IMD38

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	WB	1:1000 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS (pH 7.4), 0.02% Sodium azide and 10% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	10% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 9636 Human Swiss-port # P05161 Human
Gene Symbol	ISG15
Gene Full Name	ISG15 ubiquitin-like modifier
Background	The protein encoded by this gene is a ubiquitin-like protein that is conjugated to intracellular target proteins upon activation by interferon-alpha and interferon-beta. Several functions have been ascribed to the encoded protein, including chemotactic activity towards neutrophils, direction of ligated target proteins to intermediate filaments, cell-to-cell signaling, and antiviral activity during viral infections. While conjugates of this protein have been found to be noncovalently attached to intermediate filaments, this protein is sometimes secreted. [provided by RefSeq, Dec 2012]
Function	Ubiquitin-like protein which plays a key role in the innate immune response to viral infection either via its conjugation to a target protein (ISGylation) or via its action as a free or unconjugated protein. ISGylation involves a cascade of enzymatic reactions involving E1, E2, and E3 enzymes which catalyze the conjugation of ISG15 to a lysine residue in the target protein. Its target proteins include IFIT1, MX1/MxA, PPM1B, UBE2L6, UBA7, CHMP5, CHMP2A, CHMP4B and CHMP6. Can also isgylate: EIF2AK2/PKR which results in its activation, DDX58/RIG-I which inhibits its function in antiviral signaling response, EIF4E2 which enhances its cap structure-binding activity and translation-inhibition activity, UBE2N and UBE2E1 which negatively regulates their activity, IRF3 which inhibits its ubiquitination and degradation and FLNB which prevents its ability to interact with the upstream activators of the JNK cascade thereby inhibiting IFNA-induced JNK signaling. Exhibits antiviral activity towards both DNA and RNA viruses, including influenza A, HIV-1 and Ebola virus. Restricts HIV-1 and ebola virus via disruption of viral budding. Inhibits the ubiquitination of HIV-1 Gag and host TSG101 and disrupts their interaction, thereby preventing assembly and release of virions from infected cells. Inhibits Ebola virus budding mediated by the VP40 protein by disrupting ubiquitin ligase activity of NEDD4 and its ability to ubiquitinate VP40. ISGylates influenza A virus NS1 protein which causes a loss of function of the protein and the inhibition of virus replication. The secreted form of ISG15 can: induce natural killer cell proliferation, act as a chemotactic factor for neutrophils and act as a IFN-gamma-inducing cytokine playing an essential role in antimycobacterial immunity. [UniProt]
Calculated Mw	18 kDa
PTM	S-nitrosylation decreases its dimerization, thereby increasing the availability as well as the solubility of monomeric ISG15 for its conjugation to cellular proteins. Induced as an inactive, precursor protein that is cleaved by specific proteases to expose the C-terminal diglycine (LRLRGG) motif. This motif is essential not only for its conjugation to substrates but also for its recognition by the relevant processing proteases.

Images



ARG56931 anti-ISG15 antibody [3E5] FACS image

Flow Cytometry: HeLa cell line stained with ARG56931 anti-ISG15 antibody [3E5] at 2-5 μg for 1×10^6 cells. Secondary antibody: Goat antiMouse IgG Alexa fluor 488 conjugate.



ARG56931 anti-ISG15 antibody [3E5] WB image

Western blot: 40 µg of 1) HeLa cell lysate, 2) MCF7 cell lysate stained with ARG56931 anti-ISG15 antibody [3E5] at 1:1000.