

ARG57383
anti-ATP5L antibodyPackage: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes ATP5L
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	ATP5L
Species	Human
Immunogen	Recombinant Protein of Human ATP5L.
Conjugation	Un-conjugated
Alternate Names	ATPase subunit g; ATP5JG; ATP synthase subunit g, mitochondrial

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	ATP5L
Gene Full Name	ATP synthase, H ⁺ transporting, mitochondrial Fo complex, subunit G
Background	Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F ₁ , and the membrane-spanning component, F _o , which comprises the proton channel. The F ₁ complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The F _o seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the g subunit of the F _o complex. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Jun 2010]
Function	Mitochondrial membrane ATP synthase (F ₁ F ₀ ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F ₁ - containing the extramembraneous catalytic core, and F ₀ - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F ₁ is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F ₀ domain. Minor subunit located with subunit a in the membrane. [UniProt]
Calculated Mw	11 kDa

Images



HeLa

ARG57383 anti-ATP5L antibody WB image

Western blot: HeLa cell lysate stained with ARG57383 anti-ATP5L antibody.