

## ARG57442 anti-CIDE B antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes CIDE B
Tested Reactivity	Hu, Ms
Tested Application	IHC-P, WB
Specificity	CIDE-B antibody has no cross activity to CIDE-A.
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CIDE B
Immunogen	Synthetic peptide (15 aa) within the last 50 aa of Human CIDE-B.
Conjugation	Un-conjugated
Alternate Names	Cell death-inducing DFFA-like effector B; Cell death activator CIDE-B

### Application Instructions

Application table	Application	Dilution
	IHC-P	5 - 20 µg/ml
	WB	0.5 - 2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse small intestine tissue lysate	

### Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.02% Sodium azide
Preservative	0.02% Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	CIDEB
Gene Full Name	cell death-inducing DFFA-like effector b
Function	Activates apoptosis. [UniProt]
Calculated Mw	25 kDa


## Images

---




ARG57442 anti-CIDE B antibody IHC image

Immunohistochemistry: Human small intestine cells stained with ARG57442 anti-CIDE B antibody at 20  $\mu\text{g/ml}$  dilution.



ARG57442 anti-CIDE B antibody IHC-P image

Immunohistochemistry: CIDE-B in Human small intestine tissue stained with ARG57442 anti-CIDE B antibody at 5  $\mu\text{g/ml}$  dilution.



ARG57442 anti-CIDE B antibody WB image

Western blot: Mouse small intestine tissue lysate stained with ARG57442 anti-CIDE B antibody at 0.5, 1 and 2  $\mu\text{g/ml}$  dilution.