

**ARG57665**  
**anti-NAA40 / NatD antibody**Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes NAA40 / NatD
Tested Reactivity	Hu, Ms
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	NAA40 / NatD
Species	Human
Immunogen	Synthetic peptide (15 aa) within the last 50 aa of Human NAA40 / NatD.
Conjugation	Un-conjugated
Alternate Names	N-acetyltransferase 11; N-alpha-acetyltransferase 40; EC 2.3.1.-; PATT1; NatD catalytic subunit; NAT11

### Application Instructions

Application table	Application	Dilution
	IHC-P	5 - 20 µg/ml
	WB	1 - 2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	WB: Human thymus tissue lysate.	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.02% Sodium azide
Preservative	0.02% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	NAA40
Gene Full Name	N(alpha)-acetyltransferase 40, NatD catalytic subunit
Function	Responsible for the acetylation of the N-terminal residues of histones H4 and H2A. [UniProt]
Calculated Mw	27 kDa

## Images




ARG57665 anti-NAA40 / NatD antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse thymus tissue stained with ARG57665 anti-NAA40 / NatD antibody at 5 µg/ml dilution.



ARG57665 anti-NAA40 / NatD antibody WB image

Western blot: Human thymus tissue lysate stained with ARG57665 anti-NAA40 / NatD antibody at 1 or 2 µg/ml dilution.



ARG57665 anti-NAA40 / NatD antibody IHC image

Immunohistochemistry: Mouse thymus tissue stained with ARG57665 anti-NAA40 / NatD antibody at 20 µg/ml dilution.