

**ARG58247**  
anti-AADAC antibodyPackage: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes AADAC
Tested Reactivity	Hu, Ms
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	AADAC
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 170-399 of Human AADAC (NP_001077.2).
Conjugation	Un-conjugated
Alternate Names	EC 3.1.1.3; CES5A1; Arylacetamide deacetylase; DAC

### Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A549	
Observed Size	46 kDa	

### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

---

Gene Symbol	AADAC
Gene Full Name	arylacetamide deacetylase
Background	Microsomal arylacetamide deacetylase competes against the activity of cytosolic arylamine N-acetyltransferase, which catalyzes one of the initial biotransformation pathways for arylamine and heterocyclic amine carcinogens [provided by RefSeq, Jul 2008]
Function	Displays cellular triglyceride lipase activity in liver, increases the levels of intracellular fatty acids derived from the hydrolysis of newly formed triglyceride stores and plays a role in very low-density lipoprotein assembly. Displays serine esterase activity in liver. Deacetylates a variety of arylacetamide substrates, including xenobiotic compounds and procarcinogens, converting them to the primary arylamide compounds and increasing their toxicity. [UniProt]
Calculated Mw	46 kDa
PTM	Glycosylation is required for enzyme activity. [UniProt]
Cellular Localization	Endoplasmic reticulum membrane, Single-pass type II membrane protein, Microsome membrane. [UniProt]

## Images

---



ARG58247 anti-AADAC antibody WB image

Western blot: 25 µg of A549 cell lysate stained with ARG58247 anti-AADAC antibody at 1:1000 dilution.